arXiv:astro-ph/0108336v2 11 Sep 2001

To appear in *The Astrophysical Journal Supplement*

Sulfur, Chlorine, and Argon Abundances in Planetary Nebulae. IIA: Observations of a Southern Sample

J.B. Milingo[1]

Department of Physics, Gettysburg College, Gettysburg, PA 17325; jmilingo@gettysburg.edu

K.B. Kwitter

Department of Astronomy, Williams College, Williamstown, MA 01267;
kkwitter@williams.edu

R.B.C. Henry[1]

Department of Physics & Astronomy, University of Oklahoma, Norman, OK 73019;
henry@mail.nhn.ou.edu

and

R.E. Cohen[2]

Department of Astronomy, Wesleyan University, Middletown, CT 06459;
recohen@wesleyan.edu

ABSTRACT

In this paper we present fully reduced and dereddened emission line strengths for a sample of 45 southern Type II planetary nebulae (PNe). The spectrophotometry for these PNe covers an extended optical/near-IR range from 3600 - 9600 Å. This PN study and subsequent analysis (presented in a companion paper), together with a similar treatment for a northern PN sample, is aimed at addressing the lack of homogeneous, consistently observed, reduced, and analyzed data sets that include the near-IR [S III] lines at 9069 and 9532 Å. The use of Type II objects only is intended to select disk nebulae that are uncontaminated by nucleosynthetic products of the progenitor star. Extending spectra redward to include the strong [S III] lines enables us to look for consistency between S^{+2} abundances inferred from these lines and from the more accessible, albeit weaker, [S III] line at λ6312.

Subject headings: ISM: abundances – planetary nebulae: general – planetary nebulae: individual (Cn2-1, Fg 1, He2-21, He2-37, He2-48, He2-55, He2-115, He2-123, He2-138, He2-140, He2-141, He2-157, He2-158, IC 1297, IC 2448, IC 2501, IC 2621, IC 4776, J320, M1-5, M1-25, M1-34, M1-38, M1-50, M1-54, M1-57, M2-10, M3-4, M3-6, M3-15, NGC 2792, NGC 2867, NGC 3195, NGC 3211, NGC 3242, NGC 5307, NGC 6309, NGC 6439, NGC 6563, NGC 6565, NGC 6629, PB6, PC14, Pe1-18, Th2-A) – stars: evolution

1. Introduction

The study of the origin and buildup of the chemical elements in the universe relies heavily on the availability of accurately measured abundances for constraining model predictions. This paper is the second in a series that endeavors to measure abundances of S, Cl, and Ar, all relative to O, and to compare our empirical results with model calculations available in the literature.

One of the salient features of this project is to provide a large set of carefully and consistently measured abundances based upon a homogeneous, spectrophotometric data set which extends from 3600-9600 Å. An important asset of this extended coverage is that it includes the strong nebular lines of [S III] $\lambda\lambda$9069,9532 which have never before been used extensively to study the S abundances in planetary nebulae (PNe). Previously, S^{+2} abundances have been inferred primarily by using the auroral line of [S III] at λ6312, but this line is naturally weak and extremely temperature-sensitive, raising some doubts about the accuracy of the ion abundances derived. The inclusion of both the auroral and nebular lines of [S III] allows us to compute S^{+2} abundances from both line types and to comment on the common use of the auroral line by other investigators to study sulfur abundances.

Kwitter & Henry (2001; hereafter Paper I) observed 19 northern Type II PNe over the same spectral region as described above. The reader is referred to their paper for a more extensive discussion of the motivation for this project. The current paper, Paper IIA in the series, presents spectrophotometric measurements for 45 southern Type II PNe; the analysis of the data and calculation of abundances based upon the data will follow in Paper IIB

---

[1] Visiting Astronomer, Cerro Tololo Interamerican Observatory, National Optical Astronomy Observatories, which is operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under cooperative agreement with the National Science Foundation.

[2] Keck Northeast Astronomy Consortium Summer Fellow at Williams College, 2001

(Milingo, Henry, & Kwitter, in preparation). The current study, then, is complementary to Paper I. Future papers in the series will consider objects for which spectroscopic measurements are available in the IR from the ISO satellite along with a final interpretation of abundance patterns observed within the entire sample.

In the next section, we present the details of the observations and data reductions (§2.1), along with a careful consideration of the accuracy of the measured line strengths (§2.2). A brief summary is given in §3.

2. Data

2.1. Observations

Observations were obtained at CTIO in 1997 from 29 March - 3 April using the 1.5m telescope and cassegrain spectrograph with Loral 1K CCD. The 1200 x 800 Loral 1K CCD has 15μ pixels. We used a $5''$ x $320''$ extended slit in the E-W direction which accommodated most of the the angularly small objects in this collection. Perpendicular to dispersion the scale was $1.3''$/pixel. Gratings #22 and #9 were used to obtain extended spectral coverage from 3600-9600 Åwith overlap in the Hα region. Both gratings have nominal wavelength dispersions of 2.8 Å/pixel and 8.6 Å FWHM resolution. Table 1 lists the objects observed, their angular sizes in arcseconds, and the exposure times in seconds for the blue and red spectral regions observed with gratings #9 and #22 respectively. Most of the PNe were observed centered along the slit. For the angularly small objects, the central star was unavoidably included. In the case of a few angularly large objects we centered on the brightest portion of the nebula. Nightly calibration frames were obtained including bias exposures, twilight flats, dome flats, HeAr comparison spectra for wavelength calibration and standard star spectra for flux sensitivity calibration.

The Loral 1K CCD produces significant fringing at wavelengths beyond ∼7500 Å. Assuming less than ideal fringe removal via dome flats, reported amplitudes for interference fringes are ±1.25% at 7500 Å, ±2.25% at 8000 Å, ±5.7% at 8500 Å, ±8% at 9000 Å, and at the longest wavelengths we measure ±9.8% at 9500 Å. We note this contribution to the uncertainty in our line intensities measured at wavelengths longer than 7500 Å.

The original two-dimensional spectra were reduced and calibrated using standard long-slit spectrum reduction methods in IRAF[3]. One-dimensional spectra were extracted from the original two-dimensional images interactively using the *kpnoslit* package. Line fluxes were measured with *splot*. The final calibrated, merged spectrum of IC 2621 is shown in Fig. 1 as an illustration of our data; important emission lines are identified.

---

[3] IRAF is distributed by the National Optical Astronomy Observatories, which is operated by the As-

2.2. Line Strengths

The focus of this paper is the listing of line strengths presented in Tables 2A-H. Each table shows the line identification and wavelength in the first column. Fluxes uncorrected for reddening are presented in columns labeled F(λ), where these flux values have been normalized to Hβ=100 using our observed value of log $F_{H\beta}$ shown in the last row of the table. These line strengths in turn were corrected for reddening by assuming that the dereddened relative strength of Hα/Hβ=2.86 (Hummer & Storey 1987, N_e=100 cm^{-3}, T_e=10,000 K) and computing the logarithmic extinction quantity c shown in the penultimate line of the table. Values for the reddening coefficients, f(λ), are listed in column (2), where we employed the extinction curve of Savage & Mathis (1979). Intensities, I(λ), were obtained by multiplying F(λ), the observed ratio relative to Hβ, by dexp[cf(λ)], where dexp[y]=10^y. In general, intensities of strong lines, i.e. those whose strengths are equal to or greater than that of Hβ, have measurement uncertainties $\leq$10%; single colons indicate uncertainties of more than $\sim$25%, and double colons denote uncertainties exceeding $\sim$50%. This includes estimated errors in the c values of $\sim$0.15. Our uncertainty estimates are based on extensive previous experience with similar types of measurements. However, we also considered consistency between measurements performed independently on the data by two of us (JBM and KBK), as well as the quality of agreement between theory and observation for line ratios which have theoretically determined values. (See discussion of Table 3B below.)

As a check on our reddening correction method, we have also calculated the extinction quantity, c, based on both the Paschen-10 and Paschen-8 lines at λ9014 and λ9546, respectively. Results of this exercise are presented in Table 3A, where for each object given in the first column we list the values of c(Hα), c(P10), and c(P8). All three values were calculated based upon theoretical line ratios given in Pengelly (1964). Since c is a measure of extinction column length, in principle all three values of c for any one object should be the same.

Figure 2 is a plot of c(P10) (open circles) or c(P8) (open squares) versus c(Hα) for objects for which these values could be computed. The solid diagonal line indicates the track for the ideal one-to-one ratio. With the exception of a few outliers, agreement is

---

sociation of Universities for Research in Astronomy, Inc. (AURA) under cooperative agreement with the National Science Foundation.

satisfactory, although there is a suggestion of a systematic trend in which c for the Paschen lines tends to be slightly lower for a given c(Hα). This effect could arise if telluric absorption in the region of the Paschen lines suppresses these lines, causing the values of c derived from them to be smaller. The same effect would result if Hα were systematically overestimated due to [N II]-Hα deblending problems. It should also be noted that the deconvolution of P8 from [S III] λ9532 was difficult for most of these objects for two reasons. First, the ratio of P8/λ9532 is in many cases only a few percent. Second, this blend falls near the edge of the spectrum where the focus is poorer, leading to broadened lines which are harder to deconvolve.

An additional check for data quality is provided by information in Table 3B, where for each object we list the measured values of seven line ratios that are set solely by atomic parameters and thus are unaffected by environmental conditions. The ratios are defined in the table footnote. Values for each ratio were computed from dereddened line strengths in Tables 2. We omitted ratios where either one of the lines is marked with a double colon in Table 2. For [Ne III], our spectral resolution was insufficient to explicitly separate [Ne III] λ3968 from Hϵ; we therefore calculated the Hϵ intensity from the theoretical ratio of Hϵ/Hβ in Hummer & Storey (1987), and subtracted it from the blend. Note that the collection of intrinsically-constant ratios in Table 3B samples regions all across our spectral coverage. The penultimate line in the table gives the average for the column above it, while the last line shows the theoretical value. In a perfect world the theoretical value and all the numbers above it would agree precisely; the level of imperfection is indicated by the standard deviations associated with the means and by a general inspection of all of the numbers in any one column. There is good agreement for all line ratios.

Finally, while we do not explicitly list them here, several objects appearing in the present sample were also a part of the northern sample discussed in Paper I. A comparison of the raw line strengths and derived c(Hα) values between these two samples shows very good agreement. Given that the two sets of data were observed, reduced, and measured completely independently by different individuals using different instruments, this favorable agreement increases our confidence in both our method and in the resulting line strengths in each sample.

3. Summary

We have described spectrophotometric observations between 3600-9600 Å and presented an extensive list of line strengths for a sample of 45 southern Type II planetary nebulae. In a companion work, Paper IIB (Milingo, Henry, & Kwitter 2002) we will use these data to derive electron temperatures and densities, as well as ionic and elemental abundances. These two papers are part of a large study of PNe described in detail in Paper I, whose purpose is to produce accurate measurements of interstellar S/O, Cl/O, and Ar/O ratios and to compare these with predictions from theoretical models of stellar nucleosynthesis.

We are grateful to the CTIO TAC for granting us observing time and to the IRAF staff for their ready answers. Our research is supported by NSF grant AST-9819123.

Table 1. Object and Exposure Time Data

Object	Angular Diameter ($''$)[1]	Total Blue Exp (sec)	Total Red Exp (sec)
Cn2-1	2.4	360	720
Fg 1	16	1500	900
He2-21	2.4	1320	3600
He2-37	23	3300	600
He2-48	14	3000	900
He2-55	18	2700	1500
He2-115	3	1800	300
He2-123	4.6	900	420
He2-138	7	90	530
He2-140	2.6	780	300
He2-141	14	1500	900
He2-157	3	1200	900
He2-158	2	420	600
IC 1297	7	210	210
IC 2448	9	390	300
IC 2501	2	520	1920
IC 2621	5	210	360
IC 4776	7.5	45	405
J320	6.4	1140	840
M1-5	2	1800	540
M1-25	4.6	900	360
M1-34	11.2	1800	300
M1-38	3.3	300	360
M1-50	5.6	1020	300
M1-54	13	1200	360
M1-57	8.4	1800	300
M2-10	4	1800	300
M3-4	13.8	3600	3000
M3-6	8.2	630	1860
M3-15	4.2	2100	300
NGC 2792	13	2400	300
NGC 2867	14	390	240
NGC 3195	42	1200	900
NGC 3211	16	480	600
NGC 3242A	25	450	480
NGC 3242B	25	225	480
NGC 5307	12.5	420	420
NGC 6309	15.5	900	600
NGC 6439	5	600	360
NGC 6563	47.5	1200	600
NGC 6565	13.6	420	360
NGC 6629	15.5	420	360
PB6	11	···	2100
PC14	7	420	300
Pe1-18	6.8	2100	600
Th2-A	23	1800	1200

[1] Taken from Acker et al. (1992).

Table 2A. Line Strengths

		Cn2-1		Fg 1		He2-21		He2-37		He2-48		He2-55	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
[O II] λ 3727	0.29	20.6	33.4	30.2	36.1	7.6	10.5	255	405	213	272	13.8	20.7
He II + H10 λ 3797	0.27	2.8	4.4	4.1	4.8	3.5	4.3	2.8:	4.3:	3.6:	4.5:	···	···
He II + H9 λ 3835	0.26	3.9	6.0	4.8	5.7	4.9	5.9	3.7:	5.6:	4.6:	5.7:	13.6:	19.5:
[Ne III] λ 3869	0.25	81.3	123	55.2	64.3	60.4	72.6	121	180	83.7	103	62.7	88.9
He I + H8 λ 3889	0.25	10.8	16.2	18.5	21.5	14.2	17.0	11.6	17.1	19.1	23.5	5.7:	8.0:
Hϵ + [Ne III] λ 3968	0.22	34.4	49.8	30.3	34.7	30.5	35.9	50.1	71.4	41.7	50.4	33.7	46.0
He I + He II λ 4026	0.21	1.3:	1.9:	2.3	2.6	1.7	2.0	0.8:	1.2:	1.4::	1.7::	1.7::	2.2::
[S II] λ 4070	0.20	2.7	3.7	2.2:	2.5:	1.0	1.1	2.5	3.4	0.6::	0.8::	1.3::	1.7::
He II + Hδ λ 4101	0.19	19.2	26.2	25.9	29.0	21.1	24.2	17.9	24.1	21.7	25.4	18.4	23.9
He II λ 4198	0.16	···	···	···	···	···	···	···	···	···	···	1.6:	2.0:
C II λ 4267	0.14	0.2::	0.3::	0.8:	0.9:	0.4	0.5	0.2::	0.3::	0.2::	0.2::	···	···
Hγ λ 4340	0.12	36.7	45.1	44.8	48.3	39.9	43.7	36.8	44.5	43.0	47.5	41.1	48.8
[O III] λ 4363	0.12	8.2	10.0	3.5	3.8	13.2	14.4	20.6	24.8	9.3	10.3	9.5	11.1
He I λ 4471	0.09	4.4	5.1	5.3	5.6	2.9	3.1	2.5	2.8	3.3:	3.5:	···	···
He II λ 4540	0.07	···	···	···	···	0.8	0.9	2.1	2.4	···	···	3.0	3.3
N III λ 4640	0.05	3.4	3.6	4.6	4.7	3.0:	3.1:	···	···	···	···	···	···
He II λ 4686	0.04	4.4	4.6	14.7	15.0	27.8	28.5	62.5	66.2	16.8	17.3	97.6	103
He I + [Ar IV] λ 4711	0.03	2.0	2.2	2.5	2.5	3.6	3.7	2.8	3.0	1.7::	1.7::	8.6	9.0
[Ne IV] λ 4724	0.03	···	···	···	···	···	···	···	···	···	···	···	···
[Ar IV] λ 4740	0.02	2.8	2.9	1.6	1.7	3.6	3.7	1.5	1.5	0.7::	0.7::	3.8	4.0
Hβ λ 4861	0.00	100	100	100	100	100	100	100	100	100	100	100	100
He I λ 4921	0.00	0.9::	0.9::	1.1:	1.1:	0.6:	0.6:	···	···	0.9::	0.9::	···	···
[O III] λ 4959	-0.03	554	527	224	219	376	367	688	656	338	330	294	282
[O III] λ 5007	-0.04	1719	1605	680	663	1144	1110	2132	1996	1028	993	898	848
[N I] λ 5199	-0.09	0.2	0.2	0.7	0.7	···	···	···	···	3.0	2.8	···	···
He II λ 5411	-0.13	0.33	0.27	0.9	0.8	2.8	2.5	6.8	5.5	2.0	1.8	9.7	8.1
[Cl III] λ 5517	-0.16	0.56	0.43	0.6:	0.5:	0.2:	0.2:	2.0	1.6	1.1::	0.9::	0.7:	0.5:
[Cl III] λ 5537	-0.16	0.97	0.74	0.4:	0.3:	0.3:	0.3:	1.3	1.0	0.7::	0.6::	0.8:	0.6:
[O I] λ 5577	-0.17	0.23	0.17	0.8:	0.7:	···	···	3.0::	2.3::	5.0:	4.3:	1.6::	1.3::
[N II] λ 5755	-0.21	1.1:	0.8:	0.6	0.6	0.1::	0.1::	7.0	5.0	3.5	3.0	0.2::	0.2::
C IV λ 5801	-0.22	···	···	0.3:	0.3:	0.2:	0.2:	···	···	···	···	16.6	12.3
He I λ 5876	-0.23	26.4	18.0	17.8	15.5	15.1	12.8	11.5	8.0	15.0	12.3	5.8	4.2
He II λ 6005	-0.26	···	···	···	···	···	···	···	···	0.2::	0.1::	···	···
He II λ 6038	-0.26	···	···	···	···	···	···	···	···	0.3::	0.2::	···	···
He II λ 6074	-0.27	0.1::	0.1::	···	···	···	···	···	···	0.2::	0.2::	···	···
[Ca V] λ 6087	-0.27	···	···	···	···	···	···	···	···	···	···	···	···
[K IV] λ 6101	-0.28	0.2::	0.1::	0.1::	0.1::	0.2::	0.1::	···	···	0.2::	0.1::	···	···
He II λ 6119	-0.28	···	···	···	···	···	···	···	···	···	···	0.2::	0.2::
He II λ 6172	-0.29	···	···	···	···	···	···	0.7::	0.5::	···	···	0.3::	0.2::
[O I] λ 6300	-0.31	6.7	4.0	1.5	1.2	0.7	0.6	22.5	13.7	17.1	13.1	···	···
[S III] λ 6312	-0.32	3.5	2.1	1.0	0.9	1.2	0.9	4.8	2.9	···	···	3.5	2.3
[O I] λ 6363	-0.32	2.4	1.4	0.4	0.3	···	···	6.3	3.8	5.3	4.0	···	···
He II λ 6407	-0.33	···	···	···	···	···	···	···	···	···	···	0.8:	0.5:
[Ar IV] λ 6436	-0.34	···	···	···	···	0.2::	0.1::	···	···	···	···	···	···
[N II] λ 6548	-0.36	20.7	11.5	20.2	16.3	···	···	162	92.1	82.1	60.8	5.7	3.5
Hα λ 6563	-0.36	518	286	356	286	372	286	505	286	387	286	471	286
[N II] λ 6584	-0.36	69.2	38.0	58.1	46.6	7.5	5.7	451	254	229	169	15.0	9.1

Table 2A—Continued

		Cn2-1		Fg 1		He2-21		He2-37		He2-48		He2-55	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
He I λ 6678	-0.38	9.1	4.9	5.3	4.2	4.6	3.5	5.0	2.7	4.4	3.2	2.6	1.5
[S II] λ 6717	-0.39	4.2	2.2	6.1	4.8	0.7	0.5	46.2	25.1	18.8	13.6	3.8	2.2
[S II] λ 6731	-0.39	7.3	3.8	6.3	5.0	0.9	0.6	38.6	20.9	12.9	9.3	3.2	1.9
He II λ 6891	-0.42	···	···	···	···	···	···	0.6::	0.3::	···	···	1.1:	0.6:
[Ar V] λ 7005	-0.43	···	···	···	···	0.4	0.3	···	···	···	···	0.6::	0.3::
He I λ 7065	-0.44	19.6	9.4	2.9	2.2	6.4	4.6	2.1:	1.0:	3.0:	2.:1	2.0::	1.1::
[Ar III] λ 7135	-0.45	39.6	18.7	15.7	11.9	8.6	6.2	62.4	30.5	21.8	14.9	24.5	13.1
He II λ 7178	-0.46	···	···	···	···	0.6	0.4	···	···	···	···	2.3	1.2
[Ar IV] λ 7236	-0.47	···	···	···	···	0.2	0.1	···	···	11.2:	7.6:	···	···
[Ar IV] λ 7264	-0.47	···	···	···	···	0.2	0.1	···	···	···	···	···	···
He I λ 7281	-0.47	···	···	0.8	0.6	0.9	0.6	0.8::	0.4::	···	···	3.8::	2.0::
[O II] λ 7323	-0.48	17.8	8.1	3.1	2.3	2.3	1.6	25.8	12.1	···	···	1.4::	0.7::
[Cl IV] λ 7529	-0.51	0.64	0.27	0.3:	0.2:	0.5	0.4	···	···	···	···	1.5::	0.8::
He II λ 7591	-0.52	···	···	···	···	···	···	0.8::	0.4::	···	···	1.8::	0.9::
[S I] λ 7726	-0.54	···	···	···	···	···	···	···	···	···	···	···	···
[Ar III] λ 7751	-0.54	10.4	4.3	3.3	2.4	2.0	1.3	14.0	6.0	7.5	4.7	7.2	3.4
[Cl IV] λ 8045	-0.57	1.8	0.7	0.7:	0.5:	1.2	0.8	2.5	1.0	···	···	1.7::	0.8::
He II λ 8236	-0.59	···	···	0.6	0.4	1.1	0.7	2.6::	1.0::	···	···	4.3:	1.9:
P17 λ 8467	-0.62	1.1	0.4	0.7:	0.5:	0.6	0.4	4.0::	1.5::	3.7::	2.2::	3.8:	1.6:
P16 λ 8502	-0.62	1.6	0.6	0.6:	0.4:	0.9	0.5	···	···	···	···	2.4::	1.0::
P15 λ 8544	-0.63	1.7	0.6	0.5:	0.4:	0.9	0.6	···	···	···	···	···	···
P14 λ 8598	-0.63	1.9	0.7	1.1	0.7	1.0	0.6	1.2::	0.4::	···	···	2.6::	1.1::
P13 λ 8664	-0.64	2.1	0.7	0.9	0.6	1.5	0.9	···	···	1.2::	0.7::	2.5:	1.0:
P12 λ 8750	-0.64	2.7	0.9	1.5	1.0	2.2	1.4	3.1::	1.1::	···	···	4.1:	1.7:
P11 λ 8862	-0.65	4.7	1.6	1.7	1.1	2.7	1.7	···	···	···	···	3.9::	1.6::
P10 λ 9014	-0.67	5.2	1.8	2.0	1.4	2.7	1.6	6.4::	2.2::	···	···	7.4:	2.9:
[S III] λ 9069	-0.67	83.3	27.6	23.3	15.5	10.7	6.6	73.5	25.5	13.5:	7.7:	42.0	16.6
P9 λ 9228	-0.68	8.8	2.9	3.7	2.4	4.7	2.9	6.5	2.2	4.2::	2.4::	7.1	2.7
[S III] λ 9532	-0.70	173	54.6	73.1	47.8	21.1	12.7	222	73.6	40.1:	22.3:	83.6	31.7
P8 λ 9546	-0.70	8.8	2.8	3.2	2.1	6.0	3.6	3.6:	1.2:	11.4::	6.3::	5.2	2.0
c			0.72		0.26		0.32		0.69		0.36		0.60
log $F_{H\beta}$[a]		-11.65		-11.70		-12.12		-12.73		-12.88		-12.73	

[a] ergs cm^{-2} s^{-1} in our extracted spectra

Table 2B. Line Strengths

		He2-115		He2-123		He2-138		He2-140		He2-141		He2-157$^{\mathrm{a}}$	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
[O II] λ 3727	0.29	24.5	95.8	18.7	50.3	34.3	38.9	32.7	115	47.1	71.0	62.6	186.2
He II + H10 λ 3797	0.27	1.7	6.0	1.9:	4.7:	2.9:	3.3:	1.6	5.0	3.2	4.7	···	···
He II + H9 λ 3835	0.26	1.1:	3.7:	2.4:	5.8:	4.8:	5.4:	1.8	5.7	4.3	6.2	···	···
[Ne III] λ 3869	0.25	7.1	22.9	4.3	10.1	···	···	0.3::	0.9::	67.1	95.6	···	···
He I + H8 λ 3889	0.25	4.6	14.4	7.7	17.7	8.2	9.2	4.5	13.0	10.3	14.6	···	···
Hϵ + [Ne III] λ 3968	0.22	8.7	24.4	8.7	18.5	12.2	13.4	5.4	14.2	33.3	45.6	···	···
He I + He II λ 4026	0.21	1.0::	2.8::	1.8	3.7	···	···	···	···	1.8	2.4	···	···
[S II] λ 4070	0.20	0.9::	2.2::	0.4::	0.9::	5.8:	6.3:	1.3::	3.1::	2.1	2.8	···	···
He II + Hδ λ 4101	0.19	9.8	23.7	13.1	24.8	22.9	24.8	9.8	22.0	20.9	27.2	3.1::	6.3::
He II λ 4198	0.16	···	···	···	···	···	···	···	···	···	···	···	···
C II λ 4267	0.14	0.2::	0.4::	0.6:	0.9:	···	···	0.2::	0.4::	1.0::	1.2::	···	···
Hγ λ 4340	0.12	25.1	44.8	30.8	46.9	40.9	43.2	26.1	44.5	40.5	48.2	18.3:	29.1:
[O III] λ 4363	0.12	1.7	2.9	···	···	···	···	···	···	14.5	17.1	···	···
He I λ 4471	0.09	2.9	4.5	5.0	6.8	···	···	2.0	2.9	1.4	1.6	3.1::	4.3::
He II λ 4540	0.07	···	···	···	···	···	···	···	···	2.3	2.6	···	···
N III λ 4640	0.05	1.0::	1.3::	1.6:	1.9:	···	···	3.0:	3.7:	2.4::	2.6::	2.8::	3.3::
He II λ 4686	0.04	0.9::	1.1::	0.3:	0.4:	···	···	0.6:	0.7:	78.1	82.2	···	···
He I + [Ar IV] λ 4711	0.03	0.8::	0.9::	0.8:	0.9:	···	···	0.4:	0.5:	7.9	8.2	···	···
[Ne IV] λ 4724	0.03	···	···	···	···	···	···	···	···	···	···	···	···
[Ar IV] λ 4740	0.02	···	···	···	···	···	···	···	···	6.3	6.5	···	···
Hβ λ 4861	0.00	100	100	100	100	100	100	100	100	100	100	100	100
He I λ 4921	0.00	1.1::	1.1::	1.8:	1.8:	···	···	0.9::	0.9::	···	···	···	···
[O III] λ 4959	-0.03	209	182	63.7	57.5	···	···	15.5	13.6	430	412	66.0	59.0
[O III] λ 5007	-0.04	686	565	207	180	0.3:	0.3:	54.2	45.3	1331	1256	214	183
[N I] λ 5199	-0.09	0.9	0.6	1.7	1.3	1.4	1.4	0.8::	0.5::	0.8	0.7	···	···
He II λ 5411	-0.13	···	···	···	···	···	···	···	···	8.3	6.9	···	···
[Cl III] λ 5517	-0.16	0.3:	0.2:	0.5:	0.3:	···	···	0.5::	0.3::	1.3	1.0	1.2	0.7
[Cl III] λ 5537	-0.16	1.2:	0.6:	0.8:	0.4:	···	···	0.9::	0.4::	1.0	0.8	1.1	0.6
[O I] λ 5577	-0.17	0.9:	0.4:	0.4::	0.2::	···	···	0.3::	0.1::	0.4::	0.3::	1.0::	0.5::
[N II] λ 5755	-0.21	5.6	2.1	2.9	1.4	0.8:	0.7:	6.7	2.8	1.4	1.0	···	···
C IV λ 5801	-0.22	1.1::	0.4::	0.4::	0.2::	···	···	···	···	···	···	0.3::	0.1::
He I λ 5876	-0.23	49.9	16.9	51.2	23.4	···	···	33.0	12.2	8.5	6.2	41.4	17.5
He II λ 6005	-0.26	···	···	···	···	···	···	1.1	0.4	···	···	···	···
He II λ 6038	-0.26	0.4:	0.1:	···	···	···	···	···	···	0.3	0.2	···	···
He II λ 6074	-0.27	···	···	···	···	···	···	···	···	0.2:	0.:1	···	···
[Ca V] λ 6087	-0.27	0.1::	0.01::	···	···	···	···	···	···	···	···	···	···
[K IV] λ 6101	-0.28	0.1::	0.02::	···	···	···	···	···	···	0.7::	0.5::	···	···
He II λ 6119	-0.28	0.1::	0.03::	···	···	···	···	···	···	···	···	···	···
He II λ 6172	-0.29	···	···	···	···	···	···	···	···	···	···	···	···
[O I] λ 6300	-0.31	15.3	3.5	3.3	1.1	1.6	1.4	10.1	2.6	5.2	3.4	7.7	2.4
[S III] λ 6312	-0.32	5.0	1.2	2.0	0.7	···	···	3.5	0.9	3.8	2.4	4.6	1.4
[O I] λ 6363	-0.32	4.9	1.1	0.9::	0.3::	0.4:	0.3:	2.9	0.7	1.4	0.9	1.9:	0.6:
He II λ 6407	-0.33	···	···	···	···	···	···	···	···	···	···	···	···
[Ar IV] λ 6436	-0.34	···	···	···	···	···	···	···	···	1.7	1.1	···	···
[N II] λ 6548	-0.36	142	26.8	295	87.8	79.8	68.4	481	104	39.3	23.8	335	88.3
Hα λ 6563	-0.36	1534	286	968	286	334	286	1340	286	474	286	1095	286
[N II] λ 6584	-0.36	412	75.4	872	254	215	184	1431	300	118	70.8	970	250

Table 2B—Continued

		He2-115		He2-123		He2-138		He2-140		He2-141		He2-157[a]	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
He I λ 6678	-0.38	23.6	4.0	21.3	5.9	···	···	15.4	3.0	2.7	1.6	16.0	3.9
[S II] λ 6717	-0.39	5.9	1.0	22.4	6.0	11.9	10.1	27.6	5.3	6.9	4.0	18.7	4.4
[S II] λ 6731	-0.39	12.3	2.0	34.4	9.2	22.1	18.7	52.4	9.9	8.6	5.0	34.1	8.0
He II λ 6891	-0.42	···	···	···	···	···	···	···	···	0.7	0.4	···	···
[Ar V] λ 7005	-0.43	···	···	···	···	···	···	···	···	3.5	1.9	···	···
He I λ 7065	-0.44	62.5	7.9	26.5	5.9	···	···	28.5	4.3	2.8	1.5	28.0	5.4
[Ar III] λ 7135	-0.45	127.0	15.3	52.0	11.2	···	···	65.8	9.4	3.8	2.0	75.0	13.8
He II λ 7178	-0.46	···	···	···	···	···	···	···	···	1.4	0.8	···	···
[Ar IV] λ 7236	-0.47	3.2	0.4	2.2	0.5	···	···	2.4:	0.3:	0.4::	0.2::	···	···
[Ar IV] λ 7264	-0.47	···	···	···	···	···	···	···	···	0.3	0.1	···	···
He I λ 7281	-0.47	5.7	0.6	3.4	0.7	···	···	4.1:	0.5:	0.3	0.2	3.9:	0.7:
[O II] λ 7323	-0.48	191	20.3	9.2	1.8	2.0	1.7	64.7	8.2	9.6	4.9	89.0	14.8
[Cl IV] λ 7529	-0.51	···	···	···	···	···	···	···	···	1.2	0.6	···	···
He II λ 7591	-0.52	···	···	···	···	···	···	···	···	1.3	0.6	···	···
[S I] λ 7726	-0.54	···	···	···	···	···	···	···	···	···	···	···	···
[Ar III] λ 7751	-0.54	37.3	3.0	13.3	2.1	···	···	17.0	1.7	8.4	4.0	19.3	2.6
[Cl IV] λ 8045	-0.57	···	···	···	···	···	···	···	···	1.5	0.7	···	···
He II λ 8236	-0.59	···	···	···	···	···	···	···	···	3.5	1.5	···	···
P17 λ 8467	-0.62	5.0:	0.3:	2.3	0.3	0.4	0.3	4.2:	0.3:	0.6	0.3	···	···
P16 λ 8502	-0.62	7.6	0.4	3.1	0.4	0.8	0.6	8.2:	0.6:	0.9	0.4	···	···
P15 λ 8544	-0.63	7.3	0.4	3.9	0.5	1.0	0.8	6.6:	0.5:	0.9	0.4	4.0	0.4
P14 λ 8598	-0.63	9.2	0.5	4.0:	0.5:	0.7	0.5	7.9	0.5	1.3	0.5	3.0:	0.3:
P13 λ 8664	-0.64	12.4	0.6	4.9	0.6	1.2	0.9	9.6	0.6	1.7	0.7	3.9:	0.4:
P12 λ 8750	-0.64	15.5	0.8	6.7	0.8	1.1	0.8	14.3	0.9	2.2	0.9	4.7:	0.4:
P11 λ 8862	-0.65	22.6	1.1	8.7	0.9	2.2	1.6	19.7	1.2	2.9	1.2	10.0:	0.9:
P10 λ 9014	-0.67	24.4	1.1	12.5	1.3	2.1	1.6	16.1	0.9	3.2	1.2	7.6:	0.6:
[S III] λ 9069	-0.67	369	16.2	248	25.7	5.0	3.8	508	28.7	52.9	20.7	349	28.7
P9 λ 9228	-0.68	52.2	2.2	22.6	2.3	3.9	2.9	47.5	2.6	6.9	2.6	25.6:	2.0:
[S III] λ 9532	-0.70	382:	14.7:	559	52.4	5.2	3.9	827	41.2	71.6	26.9	437	32.3
P8 λ 9546	-0.70	52.3:	2.0:	···	···	0.7	0.5	47.0	2.3	6.8	2.6	36.0	2.6
c			2.03		1.47		0.19		1.86		0.61		1.62
log $F_{H\beta}$[b]		-12.39		-12.09		-10.70		-12.43		-12.11		-12.55	

[a] The spectrum of He2-157 contains a blue continuum and broad stellar absorption especially evident in the higher H and He lines.

[b] ergs cm^{-2} s^{-1} in our extracted spectra

Table 2C. Line Strengths

		He2-158		IC 1297		IC 2448		IC 2501		IC 2621		IC 4776	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
[O II] λ 3727	0.29	105	138	49.1	51.6	4.8::	5.1::	47.6	69.4	34.7	57.5	48.6	53.6
He II + H10 λ 3797	0.27	4.7	6.1	···	···	3.9	4.1	3.6	5.1	2.5	4.1	4.3	4.8
He II + H9 λ 3835	0.26	4.9	6.3	···	···	5.9	6.2	4.4	6.2	4.6	7.2	5.1	5.6
[Ne III] λ 3869	0.25	28.5	36.0	109	114	97.5	103	56.9	78.8	96.4	149	69.2	75.4
He I + H8 λ 3889	0.25	15.9	20.0	19.2	20.0	17.2	18.1	14.4	19.8	10.5	16.1	15.5	16.8
Hϵ + [Ne III] λ 3968	0.22	21.3	26.3	50.7	52.7	46.8	48.9	32.8	43.8	42.4	62.5	37.8	40.8
He I + He II λ 4026	0.21	2.0:	2.5:	2.1	2.2	1.6	1.7	2.0	2.7	1.9	2.7	2.3	2.5
[S II] λ 4070	0.20	2.3:	2.7:	2.8:	2.8:	1.0	1.1	2.3	3.0	5.7	8.0	3.7	3.9
He II + Hδ λ 4101	0.19	20.6	24.5	26.0	26.8	27.4	28.4	21.8	27.8	20.1	27.8	24.4	26.0
He II λ 4198	0.16	···	···	···	···	···	···	···	···	0.6::	0.7::	···	···
C II λ 4267	0.14	0.3::	0.3::	0.5:	0.5:	0.4:	0.4:	0.8	1.0	0.5:	0.6:	0.1::	0.1::
Hγ λ 4340	0.12	42.8	48.1	45.7	46.7	46.6	47.8	43.5	51.1	37.3	46.3	45.7	47.7
[O III] λ 4363	0.12	2.4	2.7	9.3	9.5	16.0	16.4	4.7	5.4	18.2	22.3	5.9	6.2
He I λ 4471	0.09	4.2	4.6	4.3	4.3	3.2	3.3	5.2	5.8	2.9	3.4	4.9	5.1
He II λ 4540	0.07	···	···	1.0	1.0	1.4	1.4	···	···	1.4	1.5	···	···
N III λ 4640	0.05	1.5::	1.6::	8.5	8.6	2.8	2.8	1.0	1.0	5.1	5.5	1.1::	1.1::
He II λ 4686	0.04	···	···	34.4	34.6	40.4	40.3	···	···	38.1	40.6	···	···
He I + [Ar IV] λ 4711	0.03	0.5::	0.5::	1.8	1.8	5.4	5.4	1.1	1.1	4.4	4.7	1.1::	1.1::
[Ne IV] λ 4724	0.03	···	···	···	···	···	···	···	···	2.0	2.1	···	···
[Ar IV] λ 4740	0.02	···	···	1.6	1.6	4.7	4.7	0.6	0.6	9.2	9.6	0.4::	0.4::
Hβ λ 4861	0.00	100	100	100	100	100	100	100	100	100	100	100	100
He I λ 4921	0.00	1.7:	1.7:	0.8::	0.8::	0.6::	0.6::	1.0::	1.0::	0.6::	0.6::	0.9::	0.9::
[O III] λ 4959	-0.03	169	164	452	450	388	386	311	299	564	535	342	338
[O III] λ 5007	-0.04	527	507	1367	1357	1183	1173	956	906	1728	1608	1043	1028
[N I] λ 5199	-0.09	1.4	1.3	0.3::	0.3::	···	···	0.6	0.5	0.9::	0.8::	0.3::	0.3::
He II λ 5411	-0.13	···	···	3.0	2.9	3.2	3.2	···	···	4.2	3.3	···	···
[Cl III] λ 5517	-0.16	0.2::	0.1::	0.7	0.7	0.2	0.2	0.4	0.3	0.5	0.4	0.4	0.4
[Cl III] λ 5537	-0.16	0.5	0.4	0.8	0.7	0.2	0.2	0.7	0.6	1.2	0.9	0.6	0.6
[O I] λ 5577	-0.17	0.6::	0.5::	0.1::	0.1::	···	···	0.1::	0.1::	0.4	0.3	···	···
[N II] λ 5755	-0.21	2.7	2.2	0.8	0.8	0.1::	0.1::	2.0	1.6	8.2	5.8	1.3	1.2
C IV λ 5801	-0.22	···	···	2.6	2.5	0.4	0.4	···	···	0.5:	0.3:	···	···
He I λ 5876	-0.23	20.3	16.4	14.6	14.0	10.2	9.7	22.4	16.7	19.9	13.3	16.9	15.6
He II λ 6005	-0.26	···	···	···	···	···	···	···	···	···	···	···	···
He II λ 6038	-0.26	···	···	0.1::	0.1::	···	···	···	···	0.2::	0.1::	···	···
He II λ 6074	-0.27	···	···	···	···	0.1::	0.1::	···	···	···	···	···	···
[Ca V] λ 6087	-0.27	···	···	···	···	···	···	···	···	0.9:	0.5:	···	···
[K IV] λ 6101	-0.28	···	···	0.1::	0.1::	0.2::	0.2::	···	···	0.6:	0.4:	···	···
He II λ 6119	-0.28	···	···	···	···	···	···	···	···	···	···	···	···
He II λ 6172	-0.29	···	···	···	···	0.1	0.1	···	···	0.2:	0.1:	···	···
[O I] λ 6300	-0.31	9.8	7.3	2.3	2.2	···	···	7.7	5.1	21.1	12.3	4.9	4.4
[S III] λ 6312	-0.32	1.3	1.0	2.3:	2.2:	0.6	0.5	1.8	1.2	6.1	3.5	2.3	2.0
[O I] λ 6363	-0.32	2.1	1.6	0.5	0.5	···	···	2.5	1.7	7.1	4.0	1.6	1.4
He II λ 6407	-0.33	···	···	···	···	0.3	0.3	···	···	0.3:	0.2 :	···	···
[Ar IV] λ 6436	-0.34	···	···	···	···	0.1::	0.1::	···	···	3.0	1.7	···	···
[N II] λ 6548	-0.36	66.7	47.8	22.3	21.0	···	···	36.8::	23.2::	115	62.0	···	···
Hα λ 6563	-0.36	400	286	304	286	307	286	455	286	533	286	323	286
[N II] λ 6584	-0.36	159	113	56.3	52.9	1.1	1.0	99.0	61.9	296	158	38.3	33.9

Table 2C—Continued

		He2-158		IC 1297		IC 2448		IC 2501		IC 2621		IC 4776	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
He I λ 6678	-0.38	6.6	4.6	4.4	4.1	3.4	3.2	6.8	4.2	6.5	3.4	4.8	4.2
[S II] λ 6717	-0.39	8.6	6.0	5.7	5.3	0.2	0.1	3.3	2.0	5.6	2.9	2.4	2.1
[S II] λ 6731	-0.39	13.1	9.1	8.5	8.0	0.1	0.1	5.9	3.6	11.0	5.6	4.2	3.7
He II λ 6891	-0.42	···	···	0.2	0.2	0.2	0.2	···	···	0.6	0.3	···	···
[Ar V] λ 7005	-0.43	0.2::	0.2::	···	···	0.1	0.1	0.1::	0.1::	7.7	3.6	0.1::	0.1::
He I λ 7065	-0.44	8.8	5.8	4.7	4.4	2.6	2.4	14.5	8.2	14.8	6.9	10.1	8.7
[Ar III] λ 7135	-0.45	14.8	9.7	18.0	16.7	5.3	4.8	25.1	14.0	66.6	30.4	16.6	14.2
He II λ 7178	-0.46	···	···	0.3:	0.3:	0.5	0.4	···	···	1.7	0.8	···	···
[Ar IV] λ 7236	-0.47	···	···	0.1::	0.1::	0.2:	0.1:	1.2	0.7	1.4	0.6	···	···
[Ar IV] λ 7264	-0.47	···	···	···	···	0.1:	0.1:	···	···	···	···	···	···
He I λ 7281	-0.47	1.4	0.9	0.6	0.6	0.4	0.4	1.2	0.7	1.4	0.6	0.8	0.7
[O II] λ 7323	-0.48	17.2	11.0	5.0	4.6	0.7	0.6	21.3	11.5	54.8	23.9	17.9	15.2
[Cl IV] λ 7529	-0.51	···	···	0.2:	0.2:	0.3	0.3	···	···	1.0	0.4	0.1::	0.1::
He II λ 7591	-0.52	···	···	0.3:	0.3:	0.5	0.4	···	···	1.1:	0.5:	···	···
[S I] λ 7726	-0.54	···	···	···	···	···	···	···	···	0.2:	0.1:	···	···
[Ar III] λ 7751	-0.54	3.5	2.1	3.8	3.5	1.3	1.2	7.0	3.5	16.4	6.5	4.0	3.3
[Cl IV] λ 8045	-0.57	0.1::	0.1::	0.6	0.6	0.7	0.6	0.2	0.1	2.4	0.9	0.2	0.2
He II λ 8236	-0.59	···	···	0.8	0.7	0.9	0.8	···	···	2.5	0.9	···	···
P17 λ 8467	-0.62	1.0::	0.6::	0.4	0.3	···	···	0.8	0.3	1.1	0.4	0.5	0.4
P16 λ 8502	-0.62	1.5::	0.9::	0.5	0.5	0.5	0.5	0.9	0.4	1.5	0.5	0.6	0.5
P15 λ 8544	-0.63	1.5:	0.8:	0.7	0.6	0.6	0.5	1.1	0.5	1.7	0.6	0.7	0.6
P14 λ 8598	-0.63	1.1:	0.6:	0.6	0.5	0.7	0.6	1.4	0.6	1.7	0.6	0.8	0.6
P13 λ 8664	-0.64	2.0:	1.1:	0.9	0.8	0.9	0.8	1.6	0.7	2.5	0.8	1.0	0.8
P12 λ 8750	-0.64	1.9:	1.1:	1.2	1.1	1.0	0.9	2.0	0.9	3.2	1.1	1.2	1.0
P11 λ 8862	-0.65	2.6:	1.4:	1.6	1.4	1.5	1.3	2.9	1.2	4.5	1.5	1.7	1.4
P10 λ 9014	-0.67	2.9:	1.5:	1.8	1.6	2.0	1.7	2.9	1.2	5.3	1.7	2.0	1.6
[S III] λ 9069	-0.67	37.2	19.9	32.6	29.1	2.9	2.5	33.7	14.2	75.2	23.6	30.6	24.4
P9 λ 9228	-0.68	5.2:	2.7:	3.4	3.0	2.6	2.3	5.2	2.2	8.4	2.6	3.1	2.5
[S III] λ 9532	-0.70	40.5	21.1	94.0	83.5	5.9	5.2	77.3	31.4	338	101	91.6	72.3
P8 λ 9546	-0.70	1.7:	0.9:	1.0:	0.9:	1.9	1.7	10.1	4.1	14.9:	4.4:	1.7:	1.3:
c			0.40		0.07		0.09		0.56		0.75		0.15
log $F_{H\beta}$[a]		-12.08		-11.13		-11.14		-10.70		-11.29		-10.78	

[a] ergs cm^{-2} s^{-1} in our extracted spectra

Table 2D. Line Strengths

		J320		M1-5		M1-25		M1-34		M1-38		M1-50	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
[O II] λ 3727	0.29	6.9	8.0	25.1	57.2	42.7	106	128	291	52.8	104	5.9	12.8
He II + H10 λ 3797	0.27	3.3	3.8	1.4::	3.0::	2.2	5.1	2.8:	6.0:	1.4:	2.5:	2.5:	5.1:
He II + H9 λ 3835	0.26	4.1	4.6	2.1:	4.5:	3.6	8.0	2.3:	4.9:	3.2:	5.8:	3.1:	6.2:
[Ne III] λ 3869	0.25	89.1	102	4.5	9.1	4.4	9.5	45.1	91.6	0.9::	1.5::	64.1	125
He I + H8 λ 3889	0.25	17.3	19.7	4.9	9.9	9.1	19.5	11.3	22.6	5.7	10.0	9.8	18.8
Hϵ + [Ne III] λ 3968	0.22	43.3	48.7	8.2	15.4	10.1	20.2	26.6	50.0	6.7	11.1	30.2	54.8
He I + He II λ 4026	0.21	1.8	2.0	1.0	1.7	1.7	3.3	2.3	4.1	···	···	1.2::	2.0::
[S II] λ 4070	0.20	0.4:	0.4:	1.1	1.8	3.3	6.0	6.9:	12.1:	4.1	6.4	1.2::	2.1::
He II + Hδ λ 4101	0.19	22.2	24.5	11.3	19.2	15.0	26.8	14.6	24.8	14.7	22.7	16.8	27.7
He II λ 4198	0.16	···	···	···	···	···	···	···	···	···	···	···	···
C II λ 4267	0.14	···	···	0.2::	0.3::	0.2::	0.4::	···	···	···	···	···	···
Hγ λ 4340	0.12	41.8	44.6	27.3	38.7	33.7	49.5	33.9	48.1	33.7	44.8	32.1	44.6
[O III] λ 4363	0.12	14.5	15.4	1.5	2.1	1.3	1.9	2.8	3.9	···	···	9.4	12.9
He I λ 4471	0.09	4.5	4.7	3.1	4.0	4.8	6.4	4.6	5.9	···	···	3.4	4.3
He II λ 4540	0.07	···	···	···	···	···	···	···	···	···	···	0.7:	0.9:
N III λ 4640	0.05	1.1:	1.1:	0.9::	1.0::	33.2	38.5	2.7:	3.0:	···	···	4.7	5.3
He II λ 4686	0.04	3.4	3.5	0.2::	0.2::	6.2::	7.0::	13.7	15.2	···	···	20.8	22.9
He I + [Ar IV] λ 4711	0.03	5.0	5.1	0.6	0.7	···	···	1.7	1.9	···	···	4.0	4.3
[Ne IV] λ 4724	0.03	···	···	···	···	···	···	···	···	···	···	···	···
[Ar IV] λ 4740	0.02	3.7	3.7	0.1::	0.1::	···	···	0.5:	0.5:	···	···	4.9	5.2
Hβ λ 4861	0.00	100	100	100	100	100	100	100	100	100	100	100	100
He I λ 4921	0.00	1.3::	1.3::	1.1:	1.1:	1.9	1.9	1.2:	1.2:	···	···	1.0:	1.0:
[O III] λ 4959	-0.03	397	391	102	93.7	170	155	252	232	···	···	586	541
[O III] λ 5007	-0.04	1220	1194	324	288	549	483	791	704	1.2:	1.1:	1823	1633
[N I] λ 5199	-0.09	0.1::	0.1::	0.5:	0.4:	1.0	0.8	12.9	10.1	1.1::	0.8::	···	···
He II λ 5411	-0.13	0.2::	0.1::	···	···	1.2	0.8	2.0:	1.4:	···	···	2.7	1.9
[Cl III] λ 5517	-0.16	0.6	0.5	0.3:	0.2:	0.6	0.4	1.9	1.2	···	···	0.6:	0.4:
[Cl III] λ 5537	-0.16	0.4	0.3	0.6:	0.4:	1.5	0.9	1.7	1.1	···	···	0.9:	0.6:
[O I] λ 5577	-0.17	0.3	0.2	1.1::	0.7::	···	···	···	···	0.6::	0.4::	0.8:	0.5:
[N II] λ 5755	-0.21	0.3:	0.3:	4.6	2.6	4.7	2.5	12.6	7.0	1.7:	0.9:	0.3:	0.2:
C IV λ 5801	-0.22	0.3	0.3	···	···	35.5	18.2	···	···	···	···	···	···
He I λ 5876	-0.23	18.9	16.8	31.6	16.5	43.1	21.1	36.9	19.2	4.3:	2.5:	25.9	14.0
He II λ 6005	-0.26	···	···	···	···	···	···	···	···	···	···	···	···
He II λ 6038	-0.26	···	···	···	···	···	···	···	···	···	···	···	···
He II λ 6074	-0.27	···	···	0.1:	0.1:	···	···	···	···	···	···	0.2::	0.1::
[Ca V] λ 6087	-0.27	···	···	···	···	···	···	···	···	···	···	···	···
[K IV] λ 6101	-0.28	0.3:	0.2:	···	···	···	···	0.4::	0.2::	···	···	0.3::	0.2::
He II λ 6119	-0.28	···	···	···	···	···	···	···	···	···	···	···	···
He II λ 6172	-0.29	···	···	···	···	···	···	···	···	···	···	···	···
[O I] λ 6300	-0.31	0.9	0.8	5.8	2.4	6.1	2.3	58.3	24.2	4.5	2.2	0.9	0.4
[S III] λ 6312	-0.32	1.3	1.1	2.0	0.8	5.0	1.9	4.2	1.7	0.5::	0.2::	3.5	1.5
[O I] λ 6363	-0.32	0.3	0.2	1.5	0.6	1.9	0.7	19.7	7.9	0.8	0.3	0.2::	0.1::
He II λ 6407	-0.33	···	···	···	···	···	···	···	···	···	···	···	···
[Ar IV] λ 6436	-0.34	···	···	···	···	···	···	···	···	···	···	···	···
[N II] λ 6548	-0.36	7.1:	5.9:	120:	43.7:	234	77.4	561	205	149	65.4	···	···
Hα λ 6563	-0.36	345	286	788	286	871	286	787	286	670	286	743	286
[N II] λ 6584	-0.36	16.4	13.6	251	90.1	652	212	1689	607	412	178	23.4	8.9

Table 2D—Continued

		J320		M1-5		M1-25		M1-34		M1-38		M1-50	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
He I λ 6678	-0.38	5.2	4.2	11.9	4.1	16.7	5.2	13.6	4.7	1.0:	0.4:	10.2	3.7
[S II] λ 6717	-0.39	0.9	0.8	3.5	1.2	17.9	5.4	161	65.6	23.6	9.4	2.2	0.8
[S II] λ 6731	-0.39	1.6	1.3	6.7	2.2	32.0	9.6	175	71.3	44.4	17.6	3.9	1.4
He II λ 6891	-0.42	···	···	···	···	···	···	···	···	···	···	···	···
[Ar V] λ 7005	-0.43	···	···	0.6::	0.2::	···	···	···	···	0.5::	0.2::	0.8::	0.3::
He I λ 7065	-0.44	6.9	5.5	32.4	9.3	29.8	7.6	12.8	3.7	0.8:	0.3:	15.5	4.8
[Ar III] λ 7135	-0.45	10.3	8.1	30.2	8.4	95.2	23.4	92.7	25.9	1.6:	0.6:	37.8	11.4
He II λ 7178	-0.46	···	···	···	···	···	···	···	···	···	···	1.3	0.4::
[Ar IV] λ 7236	-0.47	0.1::	0.1::	1.7	0.4	1.5	0.4	···	···	···	···	0.6::	0.2
[Ar IV] λ 7264	-0.47	···	···	···	···	···	···	···	···	···	···	0.5::	0.1::
He I λ 7281	-0.47	0.9:	0.7:	3.8	1.0	3.2	0.7	···	···	···	···	2.4	0.7
[O II] λ 7323	-0.48	3.0	2.4	107	27.6	43.4	9.8	32.4	8.4	14.6	4.7	8.9	2.5
[Cl IV] λ 7529	-0.51	0.5:	0.4:	···	···	···	···	···	···	···	···	1.6	0.4
He II λ 7591	-0.52	···	···	···	···	···	···	···	···	···	···	1.3	0.3
[S I] λ 7726	-0.54	···	···	···	···	···	···	···	···	···	···	···	···
[Ar III] λ 7751	-0.54	2.4	1.8	8.8	1.9	27.5	5.2	25.8	5.7	0.4::	0.1::	10.4	2.5
[Cl IV] λ 8045	-0.57	1.1	0.8	···	···	···	···	···	···	···	···	5.0	1.1
He II λ 8236	-0.59	···	···	···	···	···	···	···	···	7.8	1.9	3.1	0.6
P17 λ 8467	-0.62	0.6	0.5	2.4	0.4	2.4	0.4	···	···	1.3	0.3	1.8:	0.3:
P16 λ 8502	-0.62	0.6	0.4	3.2	0.5	3.2	0.5	···	···	1.8	0.4	2.1:	0.4:
P15 λ 8544	-0.63	0.8	0.6	3.9	0.7	3.5	0.5	···	···	1.8	0.4	2.1:	0.4:
P14 λ 8598	-0.63	1.1	0.8	4.7	0.8	3.4	0.5	···	···	1.9	0.4	2.8:	0.5:
P13 λ 8664	-0.64	1.5:	1.1:	5.3	0.9	4.8	0.7	3.7::	0.6::	2.2	0.5	3.7:	0.7:
P12 λ 8750	-0.64	1.8	1.3	7.5	1.2	6.7	0.9	6.5::	1.1::	3.5	0.8	5.6:	1.0:
P11 λ 8862	-0.65	2.7	1.9	11.5	1.8	9.0	1.2	6.8::	1.1::	5.0	1.1	6.8:	1.2:
P10 λ 9014	-0.67	2.3	1.7	2.9	0.4	10.6	1.4	8.2:	1.3:	4.0	0.9	8.5:	1.5:
[S III] λ 9069	-0.67	9.8	6.9	86.2	13.1	292	36.8	241	36.7	36.1	7.4	76.9	13.0
P9 λ 9228	-0.68	3.2	2.2	20.9	3.1	18.8	2.3	11.2:	1.7:	10.6	2.1	12.5	2.1
[S III] λ 9532	-0.70	27.3	19.0	272	38.0	929	107	743	104	71.4	13.4	247	38.7
P8 λ 9546	-0.70	4.2	2.9	30.5	4.3	12.5	1.4	24.8:	3.5:	14.3	2.7	6.0::	0.9::
c			0.23		1.22		1.34		1.22		1.03		1.15
log $F_{H\beta}$[a]		-11.51		-12.04		-12.01		-12.70		-11.98		-12.17	

[a] ergs cm^{-2} s^{-1} in our extracted spectra

Table 2E. Line Strengths

		M1-54		M1-57		M2-10		M3-4		M3-6		M3-15	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
[O II] λ 3727	0.29	134	236	75.8	254	38.3	84.5	268	346	11.8	15.0	10.5	36.8
He II + H10 λ 3797	0.27	2.9	4.8	2.0::	6.3::	1.8::	3.8::	3.7	4.7	0.6::	1.7::	···	···
He II + H9 λ 3835	0.26	3.8	6.4	···	···	2.4::	4.8::	4.4	5.5	4.0	4.9	1.6::	4.8::
[Ne III] λ 3869	0.25	74.4	121	60.6	172	3.7	7.4	115	143	38.6	47.4	20.9	60.1
He I + H8 λ 3889	0.25	14.4	23.2	6.7	18.5	10.3	20.1	18.2	22.5	16.4	20.0	5.3	15.2
Hϵ + [Ne III] λ 3968	0.22	36.6	56.5	19.7	49.9	10.2	18.7	51.7	62.8	24.5	29.4	11.2	28.5
He I + He II λ 4026	0.21	2.2	3.3	0.7::	1.6::	1.7	3.0	2.5	3.0	2.0	2.4	1.0:	2.4:
[S II] λ 4070	0.20	8.3	12.1	9.4	21.3	1.6	2.7	0.9::	1.0::	0.5::	0.6::	1.4::	3.3::
He II + Hδ λ 4101	0.19	19.5	28.0	11.5	25.0	15.3	25.5	22.1	26.0	22.1	25.7	9.8	22.0
He II λ 4198	0.16	···	···	···	···	···	···	···	···	···	···	···	···
C II λ 4267	0.14	0.4:	0.5:	···	···	0.3::	0.5::	···	···	0.3::	0.3::	0.2::	0.4::
Hγ λ 4340	0.12	37.6	47.8	26.9	45.0	33.4	46.8	45.1	50.2	42.3	46.8	25.2	41.9
[O III] λ 4363	0.12	4.8	6.0	16.3	26.6	···	···	10.8	12.0	2.0	2.2	1.9	3.1
He I λ 4471	0.09	5.2	6.1	1.7	2.4	5.4	6.9	5.5:	5.9:	5.3	5.7	3.8	5.5
He II λ 4540	0.07	0.4:	0.5:	1.1:	1.5:	···	···	1.0::	1.1::	···	···	···	···
N III λ 4640	0.05	5.3	5.8	3.3:	4.1:	0.9::	1.0::	···	···	3.6	3.7	15.3::	18.9::
He II λ 4686	0.04	16.8	18.0	44.0	51.1	···	···	31.6	32.6	1.1	1.2	3.5::	4.2::
He I + [Ar IV] λ 4711	0.03	3.3	3.5	4.9	5.5	0.5::	0.6::	2.6	2.7	0.8	0.9	1.4::	1.6::
[Ne IV] λ 4724	0.03	···	···	1.8	2.0	···	···	···	···	···	···	···	···
[Ar IV] λ 4740	0.02	2.2	2.3	7.3	8.1	···	···	1.8	1.8	0.3	0.4	0.9::	1.0::
Hβ λ 4861	0.00	100	100	100	100	100	100	100	100	100	100	100	100
He I λ 4921	0.00	1.5::	1.5::	···	···	1.6::	1.6::	0.6::	0.6::	1.1:	1.1:	1.1::	1.1::
[O III] λ 4959	-0.03	305	288	655	578	48.0	44.2	371	361	235	229	328	289
[O III] λ 5007	-0.04	941	868	2173	1829	154	138	1120	1080	714	690	1071	904
[N I] λ 5199	-0.09	13.0	11.0	5.5	3.9	1.9	1.5	4.9	4.5	···	···	0.2::	0.1::
He II λ 5411	-0.13	1.8	1.4	9.5	5.4	···	···	4.0	3.5	···	···	···	···
[Cl III] λ 5517	-0.16	1.0	0.8	2.1	1.1	0.6:	0.4:	0.9	0.8	0.3	0.3	0.8	0.4
[Cl III] λ 5537	-0.16	1.1	0.8	3.0	1.5	0.5:	0.3:	0.8:	0.7:	0.4	0.3	1.8	0.9
[O I] λ 5577	-0.17	0.3::	0.2::	0.7::	0.4::	0.8	0.5	2.8:	2.4:	···	···	0.9::	0.4::
[N II] λ 5755	-0.21	14.3	9.6	25.4	10.8	2.4	1.4	5.6	4.7	0.1::	0.1::	5.6	2.3
C IV λ 5801	-0.22	···	···	···	···	···	···	···	···	2.1	1.8	65.7	25.9
He I λ 5876	-0.23	29.1	18.6	31.8	12.2	37.0	19.8	18.5	15.1	20.3	16.8	48.4	18.3
He II λ 6005	-0.26	···	···	···	···	···	···	···	···	···	···	···	···
He II λ 6038	-0.26	···	···	···	···	···	···	···	···	···	···	···	···
He II λ 6074	-0.27	···	···	···	···	···	···	0.5::	0.4::	···	···	···	···
[Ca V] λ 6087	-0.27	···	···	10.2	3.3	···	···	···	···	···	···	···	···
[K IV] λ 6101	-0.28	···	···	1.5	0.5	···	···	···	···	0.1::	0.1::	···	···
He II λ 6119	-0.28	0.2	0.1	···	···	···	···	···	···	···	···	···	···
He II λ 6172	-0.29	0.1	0.1	···	···	···	···	···	···	···	···	···	···
[O I] λ 6300	-0.31	36.8	20.1	112	30.7	4.2	1.8	24.9	19.0	···	···	9.8	2.6
[S III] λ 6312	-0.32	3.2	1.8	29.3	7.8	1.2	0.5	0.8::	0.6::	0.8	0.7	6.4	1.7
[O I] λ 6363	-0.32	12.2	6.5	37.2	9.7	1.1	0.5	7.6	5.7	···	···	3.8	1.0
He II λ 6407	-0.33	···	···	1.2	0.3	···	···	···	···	···	···	···	···
[Ar IV] λ 6436	-0.34	0.1::	0.1::	7.0	1.7	···	···	···	···	···	···	···	···
[N II] λ 6548	-0.36	478	240	620	141	212	80.5	141	103	···	···	107	23.2
Hα λ 6563	-0.36	573	286	1268	286	758	286	391	286	383	286	1316	286
[N II] λ 6584	-0.36	1428	707	1874	416	608	227	391	286	6.6	4.9	216	47.0

Table 2E—Continued

		M1-54		M1-57		M2-10		M3-4		M3-6		M3-15	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
He I λ 6678	-0.38	10.2	4.9	14.2	2.9	14.7	5.3	5.9	4.2	6.3	4.6	22.9	4.6
[S II] λ 6717	-0.39	83.3	39.5	131	26.5	24.5	8.6	14.2	10.1	0.5	0.3	15.4	3.0
[S II] λ 6731	-0.39	110	51.9	214	42.8	33.4	11.7	11.4	8.1	0.6	0.5	26.7	5.1
He II λ 6891	-0.42	···	···	2.5	0.4	···	···	···	···	···	···	···	···
[Ar V] λ 7005	-0.43	0.3::	0.2::	18.3	3.0	···	···	···	···	···	···	···	···
He I λ 7065	-0.44	10.1	4.3	26.9	4.3	11.9	3.6	4.0	2.7	6.3	4.4	43.7	6.8
[Ar III] λ 7135	-0.45	62.0	25.9	210	32.2	35.0	10.3	31.1	21.0	25.0	17.3	120	17.8
He II λ 7178	-0.46	0.8::	0.3::	8.3	1.2	···	···	···	···	···	···	···	···
[Ar IV] λ 7236	-0.47	···	···	1.2:	0.2:	···	···	···	···	0.3::	0.2::	1.1:	0.2:
[Ar IV] λ 7264	-0.47	···	···	···	···	···	···	···	···	···	···	···	···
He I λ 7281	-0.47	1.6	0.6	2.4	0.3	···	···	0.9:	0.6:	0.9	0.6	5.7	0.8
[O II] λ 7323	-0.48	22.0	8.7	252	34.5	7.0::	1.9::	15.0	9.8	2.7	1.8	47.7	6.2
[Cl IV] λ 7529	-0.51	···	···	4.4	0.5	···	···	···	···	0.1	0.1	···	···
He II λ 7591	-0.52	···	···	···	···	···	···	···	···	···	···	···	···
[S I] λ 7726	-0.54	···	···	···	···	···	···	···	···	···	···	···	···
[Ar III] λ 7751	-0.54	15.9	5.6	57.4	6.2	9.3	2.2	7.5:	4.6:	4.9	3.2	37.9	3.9
[Cl IV] λ 8045	-0.57	1.5:	0.5:	12.0	1.1	0.9::	0.2::	···	···	0.2::	0.1::	1.6	0.1
He II λ 8236	-0.59	1.4:	0.4:	11.7	1.0	···	···	0.9	0.5	···	···	···	···
P17 λ 8467	-0.62	1.1:	0.3:	3.1:	0.2:	2.5::	0.5::	2.0	1.1	0.6	0.3	4.6:	0.3:
P16 λ 8502	-0.62	1.4:	0.4:	4.3	0.3	···	···	1.2	0.7	0.8	0.5	5.7:	0.4:
P15 λ 8544	-0.63	1.4:	0.4:	6.3:	0.5:	···	···	···	···	0.9	0.5	7.9	0.6
P14 λ 8598	-0.63	1.4:	0.4:	7.9:	0.6:	2.6:	0.5:	1.0	0.6	1.2	0.7	6.9	0.5
P13 λ 8664	-0.64	2.8	0.8	8.8:	0.6:	2.7:	0.5:	···	···	1.7	1.0	10.4:	0.7:
P12 λ 8750	-0.64	3.9	1.1	9.7:	0.7:	4.5:	0.8:	5.2	3.0	2.1	1.2	13.5	0.9
P11 λ 8862	-0.65	4.2	1.2	13.8:	0.9:	5.8:	1.0:	1.4	0.8	2.4	1.4	18.6	1.2
P10 λ 9014	-0.67	5.6	1.5	17.3:	1.1:	3.5	0.6	···	···	1.2	0.7	13.8	0.8
[S III] λ 9069	-0.67	150	41.2	690	43.2	146	23.8	13.7	7.6	32.9	19.1	489	28.6
P9 λ 9228	-0.68	8.0	2.2	37.7:	2.3:	14.4	2.3	4.6:	2.5:	4.9	2.8	40.2	2.3
[S III] λ 9532	-0.70	220	57.1	2117	118	240	36.2	40.3	21.8	74.1	42.0	1494	77.0
P8 λ 9546	-0.70	2.8::	0.7::	···	···	13.3	2.0	5.0:	2.7:	7.4	4.2	20.0	1.0
c			0.84		1.80		1.18		0.38		0.35		1.84
log $F_{H\beta}$[a]		-11.97		-12.59		-12.50		-12.61		-11.22		-12.59	

[a] ergs cm^{-2} s^{-1} in our extracted spectra

Table 2F. Line Strengths

		NGC 2792		NGC 2867		NGC 3195		NGC 3211		NGC 3242A[a]		NGC 3242B[b]	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
[O II] λ 3727	0.29	4.7:	7.4:	83.2	99.2	240	282	23.4	27.7	7.7	7.7	10.9:	11.2:
He II + H10 λ 3797	0.27	2.1:	3.2:	6.8:	8.0:	3.9	4.5	4.1	4.8	4.2	4.2	4.3	4.4
He II + H9 λ 3835	0.26	4.3	6.4	6.8:	8.0:	5.6	6.5	5.8	6.7	5.9	5.9	5.9	6.0
[Ne III] λ 3869	0.25	46.0	67.8	91.0	106	70.0	80.3	83.4	96.5	99.5	99.5	103	105
He I + H8 λ 3889	0.25	14.8	21.6	16.7	19.4	15.5	17.7	13.3	15.3	18.6	18.6	20.2	20.6
Hϵ + [Ne III] λ 3968	0.22	26.2	37.0	43.3	49.5	40.6	45.9	41.9	47.7	40.1	40.1	49.5	50.5
He I + He II λ 4026	0.21	1.0:	1.4:	1.8	2.1	2.5	2.9	1.4	1.6	2.0	2.0	2.1	2.2
[S II] λ 4070	0.20	0.8:	1.0:	3.0	3.4	7.2	8.0	1.7	1.9	1.0	1.0	1.2:	1.2:
He II + Hδ λ 4101	0.19	20.8	27.8	23.3	26.1	25.8	28.6	25.5	28.4	27.4	27.4	27.6	28.1
He II λ 4198	0.16	1.1:	1.4:	0.5::	0.6::	···	···	1.3:	1.4:	0.6::	0.6::	0.1::	0.1::
C II λ 4267	0.14	0.1::	0.1::	0.8	0.9	1.2::	1.3 ::	0.4:	0.5:	0.7	0.7	0.6	0.6
Hγ λ 4340	0.12	39.9	48.3	43.5	46.9	43.9	47.0	45.5	48.9	46.7	46.7	47.0	47.5
[O III] λ 4363	0.12	14.2	17.0	12.8	13.7	2.8	3.0	22.3	23.9	13.3	13.3	13.0	13.1
He I λ 4471	0.09	0.6	0.7	3.5	3.7	5.4	5.7	1.1	1.2	3.6	3.6	4.3	4.3
He II λ 4540	0.07	3.0	3.4	1.1	1.1	···	···	2.5	2.6	1.0	1.0	0.7:	0.7:
N III λ 4640	0.05	2.4:	2.6:	3.2	3.3	2.6:	2.7:	4.0	4.1	3.8	3.8	3.5	3.5
He II λ 4686	0.04	91.8	97.0	32.2	32.9	12.7	13.0	86.8	88.6	31.1	31.1	16.8	16.9
He I + [Ar IV] λ 4711	0.03	9.3	9.7	2.4	2.4	1.1:	1.2:	11.9	12.1	4.7	4.7	4.2	4.2
[Ne IV] λ 4724	0.03	···	···	···	···	···	···	···	···	···	···	···	···
[Ar IV] λ 4740	0.02	8.2	8.5	1.8	1.8	0.5	0.5	8.9	9.0	4.1	4.1	3.6	3.6
Hβ λ 4861	0.00	100	100	100	100	100	100	100	100	100	100	100	100
He I λ 4921	0.00	···	···	···	···	1.4:	1.4:	···	···	0.6::	0.6::	0.7::	0.7::
[O III] λ 4959	-0.03	331	316	456	448	214	211	492	484	426	426	424	423
[O III] λ 5007	-0.04	1023	960	1380	1346	643	629	1491	1456	1280	1280	1276	1271
[N I] λ 5199	-0.09	···	···	0.8	0.8	5.0	4.8	0.2::	0.2::	···	···	···	···
He II λ 5411	-0.13	10.5	8.5	3.2	2.9	1.0	0.9	7.7	7.1	2.6	2.6	1.4	1.4
[Cl III] λ 5517	-0.16	0.7	0.5	0.7	0.6	0.8	0.7	1.1	1.0	0.3	0.3	0.3	0.3
[Cl III] λ 5537	-0.16	0.6	0.5	0.8	0.7	0.7	0.6	1.0	0.9	0.3	0.3	0.3	0.3
[O I] λ 5577	-0.17	0.2::	0.2::	0.1::	0.1::	0.5::	0.5::	0.2:	0.2:	···	···	···	···
[N II] λ 5755	-0.21	0.1::	0.1::	1.5	1.3	4.6	4.1	0.3::	0.2::	0.1	0.1	0.1::	0.1::
C IV λ 5801	-0.22	0.9	0.6	0.6	0.5	···	···	···	···	0.2	0.2	···	···
He I λ 5876	-0.23	4.3	3.0	14.4	12.5	18.8	16.6	5.0	4.4	11.5	11.5	12.9	12.6
He II λ 6005	-0.26	0.1::	0.1::	0.1::	0.1::	···	···	···	···	···	···	···	···
He II λ 6038	-0.26	0.2::	0.1::	···	···	···	···	···	···	0.1:	0.1:	···	···
He II λ 6074	-0.27	0.1::	0.1::	0.1::	0.1::	···	···	···	···	0.1:	0.1:	···	···
[Ca V] λ 6087	-0.27	···	···	···	···	···	···	···	···	···	···	···	···
[K IV] λ 6101	-0.28	0.8	0.5	0.2::	0.2::	···	···	0.4::	0.3::	0.3:	0.3:	0.2::	0.2::
He II λ 6119	-0.28	0.1::	0.1::	0.1::	0.1::	···	···	···	···	0.1:	0.1:	···	···
He II λ 6172	-0.29	0.4:	0.3:	0.1::	0.1::	···	···	0.2::	0.2::	···	···	···	···
[O I] λ 6300	-0.31	···	···	8.9	7.4	14.7	12.4	···	···	···	···	···	···
[S III] λ 6312	-0.32	2.9	1.8	2.2	1.8	1.6	1.3	3.7	3.1	0.8	0.8	0.7	0.7
[O I] λ 6363	-0.32	···	···	2.9	2.4	4.9	4.1	···	···	···	···	···	···
He II λ 6407	-0.33	0.8:	0.5:	0.2:	0.1:	0.1::	0.1::	0.4::	0.3::	0.1:	0.1:	0.1::	0.1::
[Ar IV] λ 6436	-0.34	2.5	1.5	0.1::	0.1::	···	···	1.4::	1.2::	···	···	···	···
[N II] λ 6548	-0.36	3.5::	2.0::	34.8	28.1	157	129	6.4::	5.2::	···	···	···	···
Hα λ 6563	-0.36	498	286	355	286	348	286	352	286	270	270	295	286
[N II] λ 6584	-0.36	6.2::	3.5::	81.9	65.8	461	378	14.6	11.8	1.7::	1.7::	3.8:	3.7:

Table 2F—Continued

		NGC 2792		NGC 2867		NGC 3195		NGC 3211		NGC 3242A[a]		NGC 3242B[b]	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
He I λ 6678	-0.38	2.7	1.5	4.5	3.6	5.7	4.6	2.2	1.8	3.5	3.5	3.6	3.5
[S II] λ 6717	-0.39	0.8	0.4	6.7	5.3	58.0	47.0	3.2	2.5	0.3	0.3	0.4	0.4
[S II] λ 6731	-0.39	1.2	0.7	9.6	7.6	48.5	39.2	3.7	3.0	0.4	0.4	0.5	0.5
He II λ 6891	-0.42	1.5	0.8	0.3:	0.2:	0.1::	0.1::	0.8	0.6	0.2	0.2	0.2::	0.2::
[Ar V] λ 7005	-0.43	6.2::	3.2::	0.2:	0.1:	···	···	3.1	2.4	0.1	0.1	0.1	0.1
He I λ 7065	-0.44	1.6	0.8	56.7	43.5	3.2	2.5	1.4	1.1	2.9	2.9	4.5	4.3
[Ar III] λ 7135	-0.45	18.7	9.3	18.7	14.2	28.2	22.0	22.8	17.6	7.2	7.2	10.5	10.1
He II λ 7178	-0.46	2.8	1.4	0.5:	0.4:	···	···	1.5	1.1	0.3:	0.3:	0.4::	0.3::
[Ar IV] λ 7236	-0.47	0.9	0.4	0.8	0.6	0.2::	0.2::	0.6	0.4	0.2	0.2	0.4	0.4
[Ar IV] λ 7264	-0.47	0.9	0.4	···	···	···	···	0.4	0.3	0.1:	0.1:	0.2	0.2
He I λ 7281	-0.47	···	···	0.7	0.5	0.9:	0.7:	0.1	0.1	0.4	0.4	0.7	0.6
[O II] λ 7323	-0.48	1.3	0.6	12.1	9.1	7.5	5.8	2.0	1.5	1.0	1.0	1.4	1.3
[Cl IV] λ 7529	-0.51	1.6	0.7	0.3	0.3	0.1	0.1	0.8	0.6	0.4	0.4	0.5	0.5
He II λ 7591	-0.52	2.7	1.2	0.4::	0.3::	1.1::	0.8::	1.6	1.2	0.2:	0.2:	0.2::	0.2::
[S I] λ 7726	-0.54	0.5::	0.2::	···	···	···	···	···	···	···	···	···	···
[Ar III] λ 7751	-0.54	4.4	1.9	4.4	3.2	5.8	4.3	4.9	3.6	1.7	1.7	2.4	2.3
[Cl IV] λ 8045	-0.57	3.9	1.6	0.8	0.6	0.1::	0.1::	2.2	1.6	0.8	0.8	1.0	1.0
He II λ 8236	-0.59	5.3	2.1	1.0	0.7	0.3::	0.2::	2.6	1.9	0.6::	0.6::	0.6	0.6
P17 λ 8467	-0.62	0.7	0.3	0.6	0.4	1.0:	0.7:	0.4	0.3	0.3	0.3	0.4	0.4
P16 λ 8502	-0.62	1.0	0.4	0.7	0.5	0.6:	0.4:	0.5::	0.3::	0.4	0.4	0.5	0.5
P15 λ 8544	-0.63	1.3	0.5	0.7	0.5	0.8:	0.6:	0.7	0.5	0.5	0.5	0.7	0.6
P14 λ 8598	-0.63	1.3	0.5	0.9	0.6	0.8:	0.5:	0.8	0.5	0.5	0.5	0.8	0.7
P13 λ 8664	-0.64	2.0	0.7	1.3	0.9	1.1:	0.8:	1.2	0.9	0.8	0.8	1.0	0.9
P12 λ 8750	-0.64	3.0	1.1	1.5	1.0	1.8:	1.2:	1.4	1.0	1.0	1.0	1.2	1.2
P11 λ 8862	-0.65	4.5	1.6	2.3	1.6	2.4	1.7	1.9	1.3	1.3	1.3	1.7	1.6
P10 λ 9014	-0.67	4.4	1.6	2.3	1.5	2.5:	1.7:	2.5	1.7	1.4	1.4	2.0	1.8
[S III] λ 9069	-0.67	19.4	6.9	22.9	15.3	33.3	23.1	26.1	17.7	4.9	4.9	7.4	7.0
P9 λ 9228	-0.68	7.6	2.7	4.5	3.0	3.1	2.1	3.9	2.7	2.4	2.4	3.5	3.3
[S III] λ 9532	-0.70	60.8	20.7	81.1	53.3	120	81.9	68.9	46.0	16.1	16.1	20.7	19.5
P8 λ 9546	-0.70	8.2	2.8	3.2	2.1	2.8:	1.9:	4.1	2.7	2.3	2.3	3.2	3.0
c			0.67		0.26		0.24		0.25		0.00		0.04
log $F_{H\beta}$[c]		-11.76		-10.91		-11.74		-11.39		-10.46		-10.52	

[a] offset 12″ E, 9″ N of the central star

[b] offset 8″ S of the central star

[c] ergs cm^{-2} s^{-1} in our extracted spectra

Table 2G. Line Strengths

		NGC 5307		NGC 6309		NGC 6439		NGC 6563		NGC 6565		NGC 6629	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
[O II] λ 3727	0.29	7.5:	9.9:	14.2	21.9	29.6	59.2	364	424	298	386	21.2	36.2
He II + H10 λ 3797	0.27	2.9:	3.8:	2.8	4.2	2.1	4.0	···	···	3.5	4.5	2.0	3.3
He II + H9 λ 3835	0.26	4.5:	5.8:	4.1	6.1	3.3	6.1	···	···	5.1	6.4	3.0	4.9
[Ne III] λ 3869	0.25	90.7	116	65.3	94.8	69.7	127	102	116	109	136	25.7	40.8
He I + H8 λ 3889	0.25	14.2	18.0	10.7	15.4	11.0	19.7	20.7	23.5	17.7	22.0	12.2	19.2
Hϵ + [Ne III] λ 3968	0.22	35.5	44.1	33.1	46.1	29.3	49.9	49.3	55.4	49.8	60.7	18.5	27.9
He I + He II λ 4026	0.21	1.5:	1.8:	1.2	1.6	2.0	3.2	···	···	2.4	2.9	1.3:	1.9:
[S II] λ 4070	0.20	0.9:	1.1:	2.1:	2.9:	3.2	5.1	···	···	10.5	12.5	···	···
He II + Hδ λ 4101	0.19	21.5	25.8	21.9	28.9	18.1	28.3	23.3	25.7	23.4	27.6	17.6	24.8
He II λ 4198	0.16	0.5:	0.6:	0.9::	1.2::	···	···	···	···	···	···	···	···
C II λ 4267	0.14	···	···	0.3::	0.4::	0.6	0.9	···	···	0.3:	0.3 :	0.3:	0.4:
Hγ λ 4340	0.12	40.7	45.9	39.0	46.9	34.5	46.3	47.5	50.7	43.3	48.3	36.9	46.3
[O III] λ 4363	0.12	14.7	16.5	10.0	11.9	6.4	8.4	7.2	7.6	7.9	8.8	2.1	2.7
He I λ 4471	0.09	2.8	3.1	1.5	1.7	3.8	4.6	3.1:	3.3:	4.7	5.1	3.9	4.6
He II λ 4540	0.07	0.9	1.0	2.4	2.7	0.3	0.4	···	···	0.5	0.5	···	···
N III λ 4640	0.05	1.7:	1.8:	5.5	5.9	6.6	7.4	···	···	1.0::	1.0::	2.7:	2.9:
He II λ 4686	0.04	40.1	41.5	77.0	81.2	20.5	22.3	15.9	16.2	15.2	15.7	1.1:	1.2:
He I + [Ar IV] λ 4711	0.03	5.5	5.7	11.3	11.8	4.0	4.3	···	···	2.0	2.1	0.7:	0.7:
[Ne IV] λ 4724	0.03	···	···	···	···	···	···	···	···	···	···	···	···
[Ar IV] λ 4740	0.02	4.8	4.9	8.5	8.8	4.7	4.9	···	···	1.4	1.5	···	···
Hβ λ 4861	0.00	100	100	100	100	100	100	100	100	100	100	100	100
He I λ 4921	0.00	0.3::	0.3::	0.2::	0.2::	0.9::	0.9::	1.8::	1.8::	0.9::	0.9::	1.1::	1.1::
[O III] λ 4959	-0.03	416	404	352	337	457	425	321	316	398	388	232	220
[O III] λ 5007	-0.04	1285	1234	1081	1017	1422	1288	989	968	1215	1171	723	670
[N I] λ 5199	-0.09	···	···	···	···	1.0	0.8	4.8::	4.6::	5.8	5.4	···	···
He II λ 5411	-0.13	4.0	3.5	7.8	6.4	2.5	1.8	1.2::	1.1::	1.5	1.3	···	···
[Cl III] λ 5517	-0.16	0.3:	0.3:	0.6	0.5	1.1	0.7	0.5::	0.4::	1.3	1.1	0.6	0.4
[Cl III] λ 5537	-0.16	0.2:	0.2:	0.7	0.5	1.5	1.0	0.2::	0.2::	1.3	1.1	0.5	0.4
[O I] λ 5577	-0.17	···	···	···	···	0.2::	0.1::	···	···	0.4	0.3	···	···
[N II] λ 5755	-0.21	0.1::	0.1::	0.4	0.3	3.7	2.3	5.2	4.6	8.3	6.9	···	···
C IV λ 5801	-0.22	···	···	0.5:	0.4:	···	···	···	···	···	···	2.1	1.4
He I λ 5876	-0.23	11.7	9.4	9.4	6.7	29.1	16.8	16.1	14.3	18.0	14.7	23.2	15.2
He II λ 6005	-0.26	···	···	0.1::	0.1::	···	···	···	···	···	···	···	···
He II λ 6038	-0.26	···	···	0.1::	0.1::	···	···	···	···	···	···	···	···
He II λ 6074	-0.27	0.1::	0.1::	0.2::	0.1::	0.1:	0.1:	···	···	···	···	···	···
[Ca V] λ 6087	-0.27	···	···	···	···	···	···	···	···	···	···	···	···
[K IV] λ 6101	-0.28	0.4:	0.3:	0.1::	0.1::	0.5:	0.2:	···	···	···	···	···	···
He II λ 6119	-0.28	0.2:	0.1:	0.3:	0.2:	···	···	···	···	···	···	···	···
He II λ 6172	-0.29	···	···	0.4	0.3	···	···	···	···	···	···	···	···
[O I] λ 6300	-0.31	0.3::	0.2::	2.5	1.6	8.3	4.0	24.5	20.8	29.8	22.6	···	···
[S III] λ 6312	-0.32	1.3::	1.0::	2.3	1.4	5.8	2.7	···	···	3.2	2.4	0.9	0.5
[O I] λ 6363	-0.32	···	···	0.8	0.5	2.8	1.3	7.9	6.7	9.6	7.2	···	···
He II λ 6407	-0.33	0.2	0.2	0.4	0.3	···	···	···	···	···	···	···	···
[Ar IV] λ 6436	-0.34	0.2	0.2	2.2	1.4	···	···	···	···	···	···	···	···
[N II] λ 6548	-0.36	···	···	13.9	8.2	119	50.9	130	108	195	142	8.3	4.3
Hα λ 6563	-0.36	405	286	487	286	672	286	345	286	393	286	553	286
[N II] λ 6584	-0.36	4.3	3.0	39.7	23.2	299	126	372	308	572	415	21.1	10.8

Table 2G—Continued

		NGC 5307		NGC 6309		NGC 6439		NGC 6563		NGC 6565		NGC 6629	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
He I λ 6678	-0.38	4.2	2.9	5.5	3.1	10.6	4.3	4.5	3.7	5.6	4.0	8.1	4.1
[S II] λ 6717	-0.39	0.7::	0.5::	4.5	2.5	17.3	6.9	24.2	19.8	55.3	39.3	1.1	0.6
[S II] λ 6731	-0.39	1.0::	0.7::	7.1	4.0	27.9	11.1	18.6	15.2	70.4	49.9	1.4	0.7
He II λ 6891	-0.42	0.4:	0.3:	0.6	0.3	0.4:	0.2:	···	···	···	···	···	···
[Ar V] λ 7005	-0.43	0.6	0.4	2.1	1.1	1.3	0.5	···	···	···	···	···	···
He I λ 7065	-0.44	3.5	2.3	4.5	2.4	15.2	5.3	5.1	4.0	5.6	3.8	9.7	4.3
[Ar III] λ 7135	-0.45	7.7	5.0	29.4	15.0	81.5	27.8	27.1	21.4	38.8	26.0	29.1	12.7
He II λ 7178	-0.46	0.8	0.5	1.4:	0.7:	1.1::	0.4::	···	···	···	···	···	···
[Ar IV] λ 7236	-0.47	0.2::	0.1::	0.2::	0.1::	1.1	0.4	···	···	···	···	1.5:	0.6:
[Ar IV] λ 7264	-0.47	0.2::	0.1::	0.3::	0.1::	···	···	···	···	···	···	···	···
He I λ 7281	-0.47	0.5	0.3	0.6	0.3	1.8	0.6	1.4	1.1	0.8	0.5	2.0	0.8
[O II] λ 7323	-0.48	2.1	1.3	6.1	3.0	16.9	5.4	14.1	11.0	24.6	16.1	4.5	1.9
[Cl IV] λ 7529	-0.51	0.7	0.4	1.4	0.6	0.6:	0.2:	···	···	0.2::	0.1::	···	···
He II λ 7591	-0.52	0.2	0.2	1.9	0.9	···	···	···	···	···	···	···	···
[S I] λ 7726	-0.54	···	···	···	···	···	···	···	···	···	···	···	···
[Ar III] λ 7751	-0.54	1.9	1.1	6.9	3.1	21.0	5.9	6.1	4.6	8.8	5.5	···	···
[Cl IV] λ 8045	-0.57	2.0	1.1	3.4	1.5	3.4	0.9	···	···	0.3	0.2	···	···
He II λ 8236	-0.59	1.4	0.8	3.0	1.2	1.6	0.4	···	···	0.6:	0.4:	···	···
P17 λ 8467	-0.62	0.5	0.3	0.9	0.4	1.3	0.3	···	···	0.5:	0.3:	1.1	0.3
P16 λ 8502	-0.62	0.7	0.4	1.1	0.4	1.9	0.4	···	···	0.7:	0.4:	1.5	0.5
P15 λ 8544	-0.63	0.9	0.5	0.9	0.4	2.5	0.6	···	···	0.9:	0.5:	1.6	0.5
P14 λ 8598	-0.63	1.0	0.6	1.2	0.5	2.6	0.6	···	···	0.6:	0.4:	1.8	0.6
P13 λ 8664	-0.64	1.3	0.7	2.3	0.9	3.5	0.8	···	···	1.2	0.7	2.4	0.7
P12 λ 8750	-0.64	1.7	0.9	2.7	1.0	4.0	0.9	1.6:	1.1:	1.6	0.9	3.1	1.0
P11 λ 8862	-0.65	2.7	1.4	2.9	1.1	6.2	1.3	1.4:	1.0:	2.0	1.1	4.7	1.4
P10 λ 9014	-0.67	2.4	1.3	4.2	1.6	5.8	1.2	2.6:	1.8:	2.5	1.4	5.1	1.5
[S III] λ 9069	-0.67	11.0	5.8	66.6	24.7	206	42.1	21.2	15.0	77.8	43.1	33.6	9.9
P9 λ 9228	-0.68	4.4	2.3	7.3	2.7	11.5	2.3	3.2:	2.2:	5.0	2.7	8.1	2.3
[S III] λ 9532	-0.70	26.9	13.7	71.3	25.4	476	90.6	18.1	12.6	185	99.8	105	29.2
P8 λ 9546	-0.70	4.7	2.4	5.2	1.9	8.2	1.6	···	···	···	···	5.0:	1.4:
c			0.42		0.64		1.03		0.23		0.38		0.80
log $F_{H\beta}$[a]		-11.61		-11.70		-11.85		-11.75		-11.46		-11.26	

[a] ergs cm^{-2} s^{-1} in our extracted spectra

Table 2H. Line Strengths

		PB6[a]		PC14		Pe1-18		Th2-A	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
[O II] λ 3727	0.29	···	···	40.6	59.7	4.8::	20.7::	72.9	137
He II + H10 λ 3797	0.27	···	···	3.5::	5.0::	2.1	8.0	···	···
He II + H9 λ 3835	0.26	···	···	5.6:	7.9:	1.2::	4.6::	···	···
[Ne III] λ 3869	0.25	···	···	75.0	105	30.2	107	111	191
He I + H8 λ 3889	0.25	···	···	16.7	23.1	4.3	14.8	7.2	12.2
Hϵ + [Ne III] λ 3968	0.22	···	···	36.2	48.7	13.7	42.1	48.0	77.7
He I + He II λ 4026	0.21	···	···	2.1:	2.8:	···	···	2.2::	3.5::
[S II] λ 4072	0.20	···	···	1.7	2.1	2.7	7.2	···	···
He II + Hδ λ 4101	0.19	···	···	21.9	28.1	9.1	23.3	17.8:	26.6:
He II λ 4198	0.16	···	···	···	···	···	···	···	···
C II λ 4267	0.14	···	···	0.6::	0.7::	···	···	···	···
Hγ λ 4340	0.12	···	···	40.1	47.3	21.7	40.4	37.8	49.4
[O III] λ 4363	0.12	···	···	4.8	5.7	4.3	7.8	13.8	17.8
He I λ 4471	0.09	···	···	4.8	5.4	3.7	5.9	3.6::	4.4::
He II λ 4540	0.07	···	···	0.3:	0.4:	···	···	1.8	2.2
N III λ 4640	0.05	···	···	17.5::	18.6::	···	···	···	···
He II λ 4686	0.04	···	···	3.3:	3.5:	0.4::	0.5::	46.8	50.6
He I + [Ar IV] λ 4711	0.03	···	···	0.6:	0.6:	0.3::	0.3::	1.8::	1.9::
[Ne IV] λ 4724	0.03	···	···	···	···	···	···	···	···
[Ar IV] λ 4740	0.02	···	···	1.4	1.4	1.9:	2.1:	2.2::	2.3::
Hβ λ 4861	0.00	···	···	100	100	100	100	100	100
He I λ 4921	0.00	···	···	1.0::	1.0::	1.7::	1.7::	···	···
[O III] λ 4959	-0.03	···	···	422	406	439	377	556	521
[O III] λ 5007	-0.04	···	···	1306	1236	1400	1137	1719	1572
[N I] λ 5199	-0.09	···	···	···	···	1.5::	1.0::	1.1::	1.0::
He II λ 5411	-0.13	···	···	0.2::	0.2::	···	···	4.2::	3.1::
[Cl III] λ 5517	-0.16	···	···	0.5::	0.4::	0.6::	0.3::	···	···
[Cl III] λ 5537	-0.16	···	···	1.0::	0.8::	1.7::	0.8::	···	···
[O I] λ 5577	-0.17	···	···	0.2::	0.2::	2.0::	0.9::	3.2::	2.2::
[N II] λ 5755	-0.21	···	···	0.6::	0.5::	14.0	5.0	5.2::	3.3::
C IV λ 5801	-0.22	···	···	31.8	23.9	12.3::	4.2::	···	···
He I λ 5876	-0.23	···	···	22.3	16.4	70.0	22.0	18.3	11.1
He II λ 6005	-0.26	···	···	···	···	···	···	···	···
He II λ 6038	-0.26	···	···	···	···	···	···	···	···
He II λ 6074	-0.27	···	···	···	···	···	···	···	···
[Ca V] λ 6087	-0.27	···	···	0.1::	0.1::	···	···	···	···
[K IV] λ 6101	-0.28	···	···	···	···	···	···	···	···
He II λ 6119	-0.28	···	···	···	···	···	···	···	···
He II λ 6172	-0.29	···	···	···	···	···	···	···	···
[O I] λ 6300	-0.31	···	···	3.6	2.4	26.6	5.6	7.5:	3.8:
[S III] λ 6312	-0.32	···	···	2.8	1.8	15.6	3.2	8.1:	4.1:
[O I] λ 6363	-0.32	···	···	1.0	0.7	···	···	2.5	1.2
He II λ 6407	-0.33	···	···	···	···	···	···	···	···
[Ar IV] λ 6436	-0.34	0.7	···	···	···	···	···	···	···
[N II] λ 6548	-0.36	25.4	···	26.1	16.3	265	44.3	102	47.4
Hα λ 6563	-0.36	100.0	···	460	286	1730	286	619	286
[N II] λ 6584	-0.36	80.7	···	65.8	40.7	677	110	260	119

Table 2H—Continued

		PB6[a]		PC14		Pe1-18		Th2-A	
Line	f(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)	F(λ)	I(λ)
He I λ 6678	-0.38	1.0	···	6.9	4.2	35.3	5.3	6.9	3.1
[S II] λ 6717	-0.39	2.0	···	8.8	5.3	15.4	2.2	16.4	7.2
[S II] λ 6731	-0.39	2.8	···	13.3	8.0	32.0	4.6	19.8	8.6
He II λ 6891	-0.42	0.3	···	···	···	···	···	···	···
[Ar V] λ 7005	-0.43	1.7	···	···	···	···	···	···	···
He I λ 7065	-0.44	1.05	···	8.7	4.9	110	12.0	5.6:	2.2:
[Ar III] λ 7135	-0.45	5.6	···	31.7	17.4	314	32.6	64.0	24.2
He II λ 7178	-0.46	0.7	···	···	···	···	···	···	···
[Ar IV] λ 7236	-0.47	0.2	···	1.0::	0.5::	4.1	0.4	···	···
[Ar IV] λ 7264	-0.47	0.2:	···	···	···	···	···	···	···
He I λ 7281	-0.47	0.2	···	1.2	0.7	9.1	0.8	···	···
[O II] λ 7323	-0.48	2.3	···	8.0	4.2	122	11.0	14.7:	5.2:
[Cl IV] λ 7529	-0.51	0.2	···	0.3:	0.2:	2.2::	0.2::	···	···
He II λ 7591	-0.52	0.3	···	···	···	···	···	···	···
[S I] λ 7726	-0.54	0.2::	···	···	···	···	···	···	···
[Ar III] λ 7751	-0.54	1.4	···	7.4	3.6	93.2	6.3	14.9	4.7
[Cl IV] λ 8045	-0.57	0.5	···	0.8	0.4	6.4	0.4	2.8::	0.8::
He II λ 8236	-0.59	1.3	···	0.2::	0.1::	···	···	2.8::	0.8::
P17 λ 8467	-0.62	···	···	0.8	0.3	8.6	0.4	2.9	0.8
P16 λ 8502	-0.62	0.2:	···	1.0	0.5	10.6	0.5	···	···
P15 λ 8544	-0.63	0.2:	···	1.1	0.5	12.6	0.6	···	···
P14 λ 8598	-0.63	0.3:	···	1.3	0.6	13.7	0.6	5.1::	1.3::
P13 λ 8664	-0.64	0.3:	···	1.4	0.6	15.4	0.6	···	···
P12 λ 8750	-0.64	0.4:	···	1.8	0.8	22.2	0.9	6.2::	1.6::
P11 λ 8862	-0.65	0.7:	···	3.0	1.3	31.9	1.2	···	···
P10 λ 9014	-0.67	0.6:		··· 3.5	1.5	37.3	1.3	···	···
[S III] λ 9069	-0.67	7.5	···	81.0	33.5	856	30.1	98.9:	23.5:
P9 λ 9228	-0.68	1.2:	···	6.1	2.5	67.4	2.2	11.0:	2.6:
[S III] λ 9532	-0.70	22.5	···	118	46.9	3184	96.7	197:	44.1:
P8 λ 9546	-0.70	0.9:		··· 1.4:	0.5:	52.5:	1.6:	···	···
c			···		0.57		2.17		0.93
log $F_{H\beta}$[b]		-11.58		-11.78		-13.01		-12.75	
		Hα							

[a] PB6 was observed only in the red; line intensities are all relative to Hα=100, and the observed flux in Hα is shown.

[b] ergs cm^{-2} s^{-1} in our extracted spectra

Table 3A. Extinction Quantity c

Object	c(Hα)	c(P10)	c(P8)
Cn2-1	0.72	0.69	0.56
Fg 1	0.26	0.05	0.00
He2-21	0.32	0.25	0.31
He2-37	0.69	···	···
He2-48	0.36	···	···
He2-55	0.60	0.93	0.22
He2-115	2.03	1.72	1.70
He2-123	1.47	1.28	···
He2-138	0.19	0.08	0.00
He2-140	1.86	1.45	1.63
He2-141	0.61	0.37	0.39
He2-157	1.62	···	1.46
He2-158	0.40	···	···
IC 1297	0.07	0.00	···
IC 2448	0.09	0.05	0.00
IC 2501	0.56	0.30	0.65
IC 2621	0.75	0.70	0.89
IC 4776	0.15	0.05	···
J320	0.23	0.15	0.09
M1-5	1.22	0.30	1.35
M1-25	1.34	1.17	···
M1-34	1.22	0.99	1.22
M1-38	1.03	0.52	0.87
M1-50	1.15	1.02	···
M1-54	0.84	0.74	0.00
M1-57	1.80	1.49	···
M2-10	1.18	0.43	0.82
M3-4	0.38	···	0.09
M3-6	0.35	0.00	0.45
M3-15	1.85	1.34	···
NGC 2792	0.67	0.58	0.51
NGC 2867	0.26	0.15	0.00
NGC 3195	0.24	0.20	0.00
NGC 3211	0.25	0.20	0.07
NGC 3242A	0.00	0.00	0.00
NGC 3242B	0.04	0.05	0.00
NGC 5307	0.42	0.17	0.16
NGC 6309	0.64	0.55	0.22
NGC 6439	1.03	0.76	···
NGC 6563	0.23	0.23	···
NGC 6565	0.38	0.20	···
NGC 6629	0.80	0.68	···
PB6	···	···	···
PC14	0.57	0.43	···
Pe1-18	2.17	2.01	···
Th2-A	0.93	···	···

Table 3B. Line Ratios[a]

Object	[Ne III]	[O III]	[N II]	[Ar III]	P10/Hβ	P8/Hβ	[S III]
Cn 2-1	3.64	3.05	3.30	4.35	0.018	0.028	1.98
Fg 1	3.45	3.03	2.86	4.96	0.014	0.021	3.08
He2-21	3.65	3.02	···	4.77	0.016	0.036	1.92
He2-37	3.25	3.04	2.76	5.08	···	···	2.89
He2-48	3.01	3.01	2.78	3.17	···	···	2.90
He2-55	2.96	3.01	2.60	3.85	0.029	0.020	1.91
He2-115	2.73	3.10	2.81	5.10	0.011	···	···
He2-123	4.04	3.13	2.89	5.33	0.013	···	2.04
He2-138	···	···	2.69	5.53	0.016	0.005	1.03
He2-140	···	3.33	2.88	···	0.009	0.023	1.44
He2-141	3.23	3.05	2.97	···	0.012	0.026	1.30
He2-157	···	3.10	2.83	5.31	···	0.026	1.13
He2-158	3.50	3.09	2.36	4.62	···	···	1.06
IC 1297	3.10	3.02	2.52	4.77	0.016	···	2.87
IC 2448	3.12	3.04	···	4.00	0.017	0.017	2.08
IC 2501	2.83	3.03	···	4.00	0.012	0.041	2.21
IC 2621	3.21	3.01	2.55	4.68	0.017	0.044	4.28
IC 4776	3.04	3.04	···	4.30	0.016	···	2.96
J320	3.11	3.05	···	4.50	0.017	0.029	2.75
M1-5	···	3.07	···	4.42	0.004	0.043	2.90
M1-25	2.26	3.12	2.74	4.50	0.014	0.014	2.91
M1-34	2.69	3.03	2.96	4.54	0.013	0.035	2.83
M1-38	···	···	2.72	4.00	0.009	0.027	1.81
M1-50	3.22	3.02	···	4.56	0.015	···	2.98
M1-54	0.59	3.01	2.95	4.63	0.015	···	1.39
M1-57	5.07	3.16	2.95	5.19	0.011	···	2.72
M2-10	2.74	3.12	2.82	4.68	0.006	0.020	1.52
M3-4	3.06	2.99	2.78	4.71	···	0.027	2.87
M3-6	3.54	3.01	···	5.41	0.007	0.042	2.20
M3-15	7.07	3.13	2.03	4.56	0.008	0.010	2.69
NGC 2792	3.23	3.04	···	4.89	0.016	0.028	3.00
NGC 2867	3.16	3.00	2.34	4.44	0.015	0.021	3.48
NGC 3195	2.69	2.98	2.93	5.12	0.017	0.019	3.55
NGC 3211	3.04	3.01	···	4.89	0.017	0.027	2.60
NGC 3242A	4.13	3.00	···	4.24	0.014	0.023	3.29
NGC 3242B	3.05	3.00	···	4.39	0.018	0.030	2.79
NGC 5307	4.12	3.05	···	4.55	0.013	0.024	2.36
NGC 6309	3.15	3.02	2.83	4.84	0.016	0.019	1.03
NGC 6439	3.74	3.03	2.48	4.71	0.012	0.016	2.15
NGC 6563	2.95	3.06	2.85	4.65	0.018	···	0.84
NGC 6565	3.05	3.02	2.92	4.73	0.014	···	2.32
NGC 6629	3.43	3.05	2.51	···	0.015	0.014	2.95
PB6	···	···	···	···	···	···	···
PC14	3.20	3.04	2.50	4.83	0.015	···	1.40
Pe1-18	4.08	3.02	2.48	5.17	0.013	0.016	3.21
Th2-A	3.09	3.02	2.51	5.15	···	···	···
Mean	3.25 ±.69	3.05±.06	2.72±.25	4.67±.46	0.014±.004	0.025±.010	2.35±.80
Theory	3.32	2.89	2.95	4.14	0.018	0.037	2.48

[a] [Ne III]: λ3869/λ3968; [O III]: λ5007/λ4959; [N II]: λ6584/λ6548; [Ar III]: λ7135/λ7751; P10/Hβ: λ9014/λ4861; P8/Hβ: λ9546/λ4861; [S III]: λ9532/λ9069.

REFERENCES

Acker, A., Ochsenbein, F., Stenholm, B., Tylenda, R., Marcout, J., & Schohn, C. 1992, The Strasbourg-ESO Catalogue of Galactic Planetary Nebulae (Garching: ESO)

Hummer, D.G., & Storey, P.J. 1987, MNRAS, 224, 801

Kwitter, K.B., & Henry, R.B.C. 2001, ApJ, *in press*; Paper I)

Pengelly R.M. 1964, MNRAS, 127, 145

Savage, B.D., & Mathis, J.S. 1979, ARA&A, 17, 73

This preprint was prepared with the AAS LaTeX macros v5.0.



Fig. 1.— Spectrum of IC 2621.



Fig. 2.— A comparison of logarithmic extinction *c* as determined using the Paschen 8 and 10 lines versus the value inferred from using Hα. The solid line shows the track for a one-to-one correspondence.